

Publications

Division of New Materials Science

Takigawa group

We have been performing nuclear magnetic resonance experiments on various quantum spin systems and strongly correlated electron systems to explore novel quantum phases with exotic ordering and fluctuation phenomena. The major achievements in the year 2015 include: (1) Investigation by ^{31}P -NMR of the quasi two dimensional antiferromagnet $\text{RbMoOPO}_4\text{Cl}$ with the frustrating nearest neighbor and the next nearest neighbor interactions that lead to the identification of a stripe type antiferromagnetic order. (2) Combined studies of ^7Li -NMR and neutron scattering experiments on the breathing pyrochlore antiferromagnet $\text{Li}(\text{Ga},\text{In})\text{Cr}_4\text{O}_8$, in particular the observation of a second order phase transition in the In 5% doped sample. (3) Investigation of spin dynamics in the quantum spin ice compound $\text{Pr}_2\text{Zr}_2\text{O}_7$ by ^{91}Zr -NQR/NMR that revealed contrasting behavior when a magnetic field is applied along $\langle 001 \rangle$ and $\langle 111 \rangle$.

1. † Structural anomalies and short-range magnetic correlations in the orbitally degenerate system Sr_2VO_4 : I. Yamauchi, K. Nawa, M. Hiraishi, M. Miyazaki, A. Koda, K. M. Kojima, R. Kadono, H. Nakao, R. Kumai, Y. Murakami, H. Ueda, K. Yoshimura and M. Takigawa, *Phys. Rev. B* **92** (2015) 064408(1-7).
2. * One-Third Magnetization Plateau with a Preceding Novel Phase in Volborthite: H. Ishikawa, M. Yoshida, K. Nawa, M. Jeong, S. Kramer, M. Horvatic, C. Berthier, M. Takigawa, M. Akaki, A. Miyake, M. Tokunaga, K. Kindo, J. Yamaura, Y. Okamoto and Z. Hiroi, *Phys. Rev. Lett.* **114** (2015) 227202(1-5).
3. † Real Space Imaging of Spin Polarons in Zn-Doped $\text{SrCu}_2(\text{BO}_3)_2$: M. Yoshida, H. Kobayashi, I. Yamauchi, M. Takigawa, S. Capponi, D. Poilblanc, F. Mila, K. Kudo, Y. Koike and N. Kobayashi, *Phys. Rev. Lett.* **114** (2015) 056402 (1-5).
4. * Single crystal ^{27}Al -NMR study of the cubic Γ_3 ground doublet system $\text{PrTi}_2\text{Al}_{20}$: T. Taniguchi, M. Yoshida, H. Takeda, M. Takigawa, M. Tsujimoto, A. Sakai, Y. Matsumoto and S. Nakatsuji, *J. Phys.: Conf. Ser.* **683** (2016) 012016(1-9).
5. * Site-selective ^{11}B NMR studies on YbAlB_4 : S. Takano, M. S. Grbic, K. Kimura, M. Yoshida, M. Takigawa, E. C. T. O. Farrell, K. Kuga, S. Nakatsuji and H. Harima, *J. Phys.: Conf. Ser.* **683** (2016) 012008(1-6).

Sakakibara group

We study magnetism and superconductivity of materials having low characteristic temperatures. These include heavy-electron systems, quantum spin systems and frustrated spin systems. The followings are some selected achievements in the fiscal year 2015. (1) Field and temperature variations of the specific heat $C(H,T)$ of the heavy fermion superconductor URu_2Si_2 ($T_c=1.4$ K) were examined at temperatures down to 200 mK. The occurrence of quasiparticle excitations due to the Doppler-shift effect was detected regardless of the field direction in $C(H)$, implying the presence of a line node. Furthermore, the polar-angle-dependence of the specific heat $C(\theta)$ under a rotating magnetic field within the ac plane exhibits a shoulder-like anomaly at $\theta \sim 45^\circ$ and a sharp dip at $\theta = 90^\circ$ ($H \parallel a$) in the moderate-field region. These features are supported by theoretical analyses based on microscopic calculations assuming the gap symmetry of $k_z(k_x+ik_y)$, whose gap structure is characterized by a combination of a horizontal line node at the equator and point nodes at the poles. The present results have settled the previous controversy over the gap structure of URu_2Si_2 and have authenticated its chiral d -wave superconductivity. (2) We examined low-temperature magnetic properties of a new verdazyl radical crystal α -2-Cl-4-F-V. Molecular orbital calculations predict that this material is a spin-1/2 quasi-one-dimensional antiferromagnet. The magnetization curve at 0.1 K shows gapless behavior like conventional one-dimensional quantum spin systems and saturates at about 5 T. A peak is observed in the temperature dependence of the heat capacity at 0 T, indicating that a three-dimensional ordering occurs at about 0.2 K. On the other hand, the temperature dependence of the magnetization at various magnetic fields shows an anomaly similar to that observed in many of gapped spin systems. This behavior is attributed to the highly frustrated nature of the interchain interactions.

1. * Field Evolution of Quantum Critical and Heavy Fermi-Liquid Components in the Magnetization of the Mixed Valence Compound β - YbAlB_4 : Y. Matsumoto, K. Kuga, Y. Karaki, Y. Shimura, T. Sakakibara, M. Tokunaga, K. Kindo and S. Nakatsuji, *J. Phys. Soc. Jpn.* **84** (2015) 024710(1-7).

* Joint research among groups within ISSP.

2. [†] Observation of a New Ordered Phase in the Kondo Semiconductor CeOs₄Sb₁₂: T. Tayama, W. Ohmachi, M. Wansawa, D. Yutani, T. Sakakibara, H. Sugawara and H. Sato, *J. Phys. Soc. Jpn.* **84** (2015) 104701(1-6).
3. ^{†*} Pauli-limited superconductivity and antiferromagnetism in the heavy-fermion compound CeCo(In_{1-x}Zn_x)₅: M. Yokoyama, H. Mashiko, R. Otaka, Y. Sakon, K. Fujimura, K. Tenya, A. Kondo, K. Kindo, Y. Ikeda, H. Yoshizawa, Y. Shimizu, Y. Kono and T. Sakakibara, *Phys. Rev. B* **92** (2015) 184509(9).
4. [†] Quantum phase near the saturation field in the S = 1/2 frustrated spin ladder: H. Yamaguchi, H. Miyagai, Y. Kono, S. Kittaka, T. Sakakibara, K. Iwase, T. Ono, T. Shimokawa and Y. Hosokoshi, *Phys. Rev. B* **91** (2015) 125104(1-5).
5. [†] S = 1/2 ferromagnetic-antiferromagnetic alternating Heisenberg chain in a zinc-verdazyl complex: H. Yamaguchi, Y. Shinpuku, T. Shimokawa, K. Iwase, T. Ono, Y. Kono, S. Kittaka, T. Sakakibara and Y. Hosokoshi, *Phys. Rev. B* **91** (2015) 085117(1-6).
6. ^{*} Unstable spin-ice order in the stuffed metallic pyrochlore Pr_{2+x}Ir_{2-x}O_{7-δ}: D. E. MacLaughlin, O. O. Bernal, L. Shu, J. Ishikawa, Y. Matsumoto, J. -J. Wen, M. Mourgil, C. Stock, G. Ehlers, C. L. Broholm, Y. Machida, K. Kimura, S. Nakatsuji, Y. Shimura and T. Sakakibara, *Phys. Rev. B* **92** (2015) 054432(1-12).
7. Field-Induced Quantum Criticality and Universal Temperature Dependence of the Magnetization of a Spin-1/2 Heisenberg Chain: Y. Kono, T. Sakakibara, C. P. Aoyama, C. Hotta, M. M. Turnbull, C. P. Landee and Y. Takano, *Phys. Rev. Lett.* **114** (2015) 037202(1-5).
8. Field-Orientation Dependence of Low-Energy Quasiparticle Excitations in the Heavy-Electron Superconductor UBe₁₃: Y. Shimizu, S. Kittaka, T. Sakakibara, Y. Haga, E. Yamamoto, H. Amitsuka, Y. Tsutsumi and K. Machida, *Phys. Rev. Lett.* **114** (2015) 147002(1-5).
9. ^{†*} Antiferromagnetic Transition in a Novel Star-Shaped High-Spin Fe(III) Tetranuclear Cluster from a Mononuclear Coordination Anion Featuring π -Extended Schiff Base Ligands: K. Takahashi, K. Kawamukai, T. Mochida, T. Sakurai, H. Ohta, T. Yamamoto, Y. Einaga, H. Mori, Y. Shimura, T. Sakakibara, T. Fujisawa, A. Yamaguchi and A. Sumiyama, *Chem. Lett.* **44** (2015) 840-842.
10. ^{*} Antiferromagnetic transition of the caged compound TmTi₂Al₂₀: N. Kase, Y. Shimura, S. Kittaka, T. Sakakibara, S. Nakatsuji, T. Nakano, N. Takeda and J. Akimitsu, *J. Phys.: Conf. Ser.* **592** (2015) 012052(1-5).
11. [†] Experimental Realization of a Quantum Pentagonal Lattice: H. Yamaguchi, T. Okubo, S. Kittaka, T. Sakakibara, K. Araki, K. Iwase, N. Amaya, T. Ono and Y. Hosokoshi, *Sci. Rep.* **5** (2015) 15327(1-6).
12. [†] First-order ferromagnetic transition of quantum spin ice system Yb₂Ti₂O₇: Y. Yasui, N. Hamachi, Y. Kono, S. Kittaka and T. Sakakibara, *SPIN* **5** (2015) 154002(1-6).
13. Low temperature magnetic properties of a new quasi-one-dimensional organic magnet α -2-Cl-4-F-V: Y. Kono, S. Kittaka, T. Sakakibara, H. Yamaguchi and Y. Hosokoshi, *Phys. Procedia* **75** (2015) 679-686.
14. Evidence for chiral d-wave superconductivity in URu₂Si₂ from the field-angle variation of its specific heat: S. Kittaka, Y. Shimizu, T. Sakakibara, Y. Haga, E. Yamamoto, Y. Onuki, Y. Tsutsumi, T. Nomoto, H. Ikeda and K. Machida, *J. Phys. Soc. Jpn.* **85** (2016) 033704(1-4).
15. Field-induced phase transitions and magnetoferroelectricity in the perfect triangular lattice antiferromagnet RbFe(MoO₄)₂ in a vertical magnetic field: H. Mitamura, R. Watanuki, N. Onozaki, Y. Amou, Y. Kono, S. Kittaka, Y. Shimura, I. Yamamoto, K. Suzuki and T. Sakakibara, *J. Magn. Magn. Mater.* **400** (2016) 70-72.
16. First-order superconducting transition of Sr₂RuO₄ investigated by magnetization and magnetic torque: S. Kittaka, A. Kasahara, T. Sakakibara, D. Shibata, S. Yonezawa, Y. Maeno, K. Tenya and K. Machida, *J. Magn. Magn. Mater.* **400** (2016) 81-83.
17. Heat capacity measurements on UBe₁₃ in rotated magnetic fields: anisotropic response in the normal state and the absence of nodal quasiparticles: Y. Shimizu, S. Kittaka, T. Sakakibara, Y. Haga, E. Yamamoto, H. Amitsuka, Y. Tsutsumi and K. Machida, *J. Magn. Magn. Mater.* **400** (2016) 52-55.
18. [†] Unconventional S = 2 alternating chain realized by a metal-radical hybrid-spin approach: H. Yamaguchi, Y. Shinpuku, Y. Kono, S. Kittaka, T. Sakakibara, M. Hagiwara, T. Kawakami, K. Iwase, T. Ono and Y. Hosokoshi, *Phys. Rev. B* **93** (2016) 115145(1-7).
19. [†] Quadrupole Order in the Frustrated Pyrochlore Tb_{2+x}Ti_{2-x}O_{7+y}: H. Takatsu, S. Onoda, S. Kittaka, A. Kasahara, Y. Kono, T. Sakakibara, Y. Kato, B. Fåk, J. Ollivier, J. W. Lynn, T. Taniguchi, M. Wakita and H. Kadowaki, *Phys. Rev. Lett.* **116** (2016) 217201(1-6).

[†] Joint research with outside partners.

20. [†] Possible observation of highly itinerant quantum magnetic monopoles in the frustrated pyrochlore Yb₂Ti₂O₇: Y. Tokiwa, T. Yamashita, M. Udagawa, S. Kittaka, T. Sakakibara, D. Terazawa, Y. Shimoyama, T. Terashima, Y. Yasui, T. Shibauchi and Y. Matsuda, Nat. Commun. **7** (2016) 10807(1-6).
21. [†] Quadrupole order in the frustrated pyrochlore magnet Tb₂Ti₂O₇: H. Takatsu, T. Taniguchi, S. Kittaka, T. Sakakibara and H. Kadowaki, J. Phys.: Conf. Series **683** (2016) 012022(1-6).
22. ^{*} Low-Energy Excitations and Ground State Selection in Quantum Breathing Pyrochlore Antiferromagnet Ba₃Yb₂Zn₅O₁₁: T. Haku, K. Kimura, Y. Matsumoto, M. Soda, M. Sera, D. Yu, R. A. Mole, T. Takeuchi, S. Nakatsuji, Y. Kono, T. Sakakibara, L. -J. Chang and T. Masuda, Phys. Rev. B (2016), in print.

Mori group

We have successfully developed and unveiled unprecedented functional properties for the molecular materials. The major achievements in 2015 are (1) to discover the peculiar hydrogen-bond-dynamics-based switching of conductivity and magnetism triggered by deuterium and charge transfers in the hydrogen-bond-unit for the purely organic conductor κ -D₃(Cat-EDT-ST)₂, (2) to develop the novel donor-acceptor-type molecular dyad with small HOMO-LUMO gap, and (3) to develop the chiral molecular conductors composed of chiral BEDT-TTF derivatives with hydrogen bonds, α' -[(R,R)-BEDT-TTF(CH₂OH)₂]₂ClO₄(H₂O). The introduction of a large variety of molecule degree of freedom to solid promises the development of new trends in functional molecular materials.

1. Dynamics of Charge Ordering in the Nonlinear Regime of θ -(BEDT-TTF)₂CsZn(SCN)₄: M. Abdel-Jawad, I. Terasaki, T. Mori and H. Mori, J. Phys. Soc. Jpn. **84** (2015) 033707(1-4).
2. [†] Phase-change memory function of correlated electrons in organic conductors: H. Oike, F. Kagawa, N. Ogawa, A. Ueda, H. Mori, M. Kawasaki and Y. Tokura, Phys. Rev. B **91** (2015) 041101(R)(1-4).
3. ^{†*} Antiferromagnetic Transition in a Novel Star-Shaped High-Spin Fe(III) Tetranuclear Cluster from a Mononuclear Coordination Anion Featuring π -Extended Schiff Base Ligands: K. Takahashi, K. Kawamukai, T. Mochida, T. Sakurai, H. Ohta, T. Yamamoto, Y. Einaga, H. Mori, Y. Shimura, T. Sakakibara, T. Fujisawa, A. Yamaguchi and A. Sumiyama, Chem. Lett. **44** (2015) 840-842.
4. Unexpected Formation of *ortho*-Benzoquinone-fused Tetraselenafulvalene (TSF): Synthesis, Structures, and Properties of a Novel TSF-based Donor-Acceptor Dyad: A. Ueda, H. Kamo and H. Mori, Chem. Lett. **44** (2015) 1538-1540.
5. Modulation of a Molecular π -Electron System in a Purely Organic Conductor that Shows Hydrogen-Bond-Dynamics-Based Switching of Conductivity and Magnetism: A. Ueda, A. Hatakeyama, M. Enomoto, R. Kumai, Y. Murakami and H. Mori, Chem. - Eur. J. **21** (2015) 15020-15028.
6. Synthesis of racemic and chiral BEDT-TTF derivatives possessing hydroxy groups and their achiral and chiral charge transfer complexes: S. J. Krivickas, C. Hashimoto, J. Yoshida, A. Ueda, K. Takahashi, J. D. Wallis and H. Mori, Beilstein J. Org. Chem. **11** (2015) 1561-1569.
7. Novel electronic ferroelectricity in an organic charge-order insulator investigated with terahertz-pump optical-probe spectroscopy: H. Yamakawa, T. Miyamoto, T. Morimoto, H. Yada, Y. Kinoshita, M. Sotome, N. Kida, K. Yamamoto, K. Iwano, Y. Matsumoto, S. Watanabe, Y. Shimoi, M. Suda, H. M. Yamamoto, H. Mori and H. Okamoto, Sci. Rep. **6** (2016) 20571(1-10).
8. 有機結晶における新機能物性開拓 — π 電子と水素の協奏—: 森 初果, Mol. Sci. **9** (2015) A0081.

Nakatsuji group

Our group explores novel quantum phases and phase transitions in rare-earth and transition metal based compounds. The followings are some relevant results obtained in 2015. (1) We discovered the first example of an antiferromagnet that exhibits the anomalous Hall effect at room temperature, the chiral antiferromagnet Mn₃Sn (2) The quantum criticality at ambient pressure in β -YbAlB₄ is found robust against pressure, and possibly forms a strange metal phase. In addition, this phase is separated from a magnetic criticality by a Fermi liquid phase stabilized under pressure (3) A field induced quantum metal-insulator transition was found in the pyrochlore iridate Nd₂Ir₂O₇, when the field is applied only along the narrow angle range close to [100]. (4) A Fermi node at the quadratic band touching in Pr₂Ir₂O₇ was found and indicates that Pr₂Ir₂O₇ should be viewed as a mother compound for various topological phases in the correlated electron system, such as Weyl semimetals, and topological insulators.

* Joint research among groups within ISSP.

1. Conduction electron spin resonance in the α -Yb_{1-x}Fe_xAlB₄ ($0 \leq x \leq 0.50$) and α -LuAlB₄ compounds: L. Holanda, G. Lesseux, E. Magnavita, R. Ribeiro, S. Nakatsuji, K. Kuga, Z. Fisk, S. Oseroff, R. Urbano, C. Rettori and P. Pagliuso, *J. Phys.: Condens. Matter* **27** (2015) 255601(1-5).
2. *Field Evolution of Quantum Critical and Heavy Fermi-Liquid Components in the Magnetization of the Mixed Valence Compound β -YbAlB₄: Y. Matsumoto, K. Kuga, Y. Karaki, Y. Shimura, T. Sakakibara, M. Tokunaga, K. Kindo and S. Nakatsuji, *J. Phys. Soc. Jpn.* **84** (2015) 024710(1-7).
3. †*Magnetic Order in the Frustrated Ising Quasi-One Dimensional Compound NaCo(acac)₃·Benzene: Y. Karaki, K. Kuga, K. Kimura, S. Nakatsuji, K. Matsubayashi and Y. Uwatoko, *J. Phys. Soc. Jpn.* **84** (2015) 084708(1-5).
4. Magnetization Anomaly due to the Non-Coplanar Spin Structure in NiS₂: T. Higo and S. Nakatsuji, *J. Phys. Soc. Jpn.* **84** (2015) 053702(1-5).
5. *X-ray Absorption Spectroscopy in the Heavy Fermion Compound α -YbAlB₄ at High Magnetic Fields: T. T. Terashima, Y. H. Matsuda, K. Kuga, S. Suzuki, Y. Matsumoto, S. Nakatsuji, A. Kondo, K. Kindo, N. Kawamura, M. Mizumaki and T. Inami, *J. Phys. Soc. Jpn.* **84** (2015) 114715(1-4).
6. Field-induced quadrupolar quantum criticality in PrV₂Al₂₀: Y. Shimura, M. Tsujimoto, B. Zeng, L. Balicas, A. Sakai and S. Nakatsuji, *Phys. Rev. B* **91** (2015) 241102(1-5).
7. Large trigonal-field effect on spin-orbit coupled states in a pyrochlore iridate: D. Uematsu, H. Sagayama, T. Arima, J. J. Ishikawa, S. Nakatsuji, H. Takagi, M. Yoshida, J. Mizuki and K. Ishii, *Phys. Rev. B* **92** (2015) 094405.
8. Optical evidence for a Weyl semimetal state in pyrochlore Eu₂Ir₂O₇: A. B. Sushkov, J. B. Hofmann, G. S. Jenkins, J. Ishikawa, S. Nakatsuji, S. Das Sarma and H. D. Drew, *Phys. Rev. B* **92** (2015) 241108(1-4).
9. *Unstable spin-ice order in the stuffed metallic pyrochlore Pr_{2+x}Ir_{2-x}O_{7- δ} : D. E. MacLaughlin, O. O. Bernal, L. Shu, J. Ishikawa, Y. Matsumoto, J. -J. Wen, M. Mourgil, C. Stock, G. Ehlers, C. L. Broholm, Y. Machida, K. Kimura, S. Nakatsuji, Y. Shimura and T. Sakakibara, *Phys. Rev. B* **92** (2015) 054432(1-12).
10. Spin Fluctuations from Hertz to Terahertz on a Triangular Lattice: Y. Nambu, J. S. Gardner, D. E. MacLaughlin, C. Stock, H. Endo, S. Jonas, T. J. Sato, S. Nakatsuji and C. Broholm, *Phys. Rev. Lett.* **115** (2015) 127202(1-5).
11. High-Field Multi-Frequency ESR in the Rare-Earth Spinel Compound CdYb₂S₄: D. Yoshizawa, T. Kida, S. Nakatsuji, K. Iritani, M. Halim, T. Takeuchi and M. Hagiwara, *Appl. Magn. Reson.* **46** (2015) 0937-9347.
12. Large anomalous Hall effect in a non-collinear antiferromagnet at room temperature: S. Nakatsuji, N. Kiyohara and T. Higo, *Nature* **527** (2015) 212-215.
13. 電子軌道の量子ゆらぎによる新しい超伝導: 松本 洋介, *パリティ* **30-7** (2015) 2-6.
14. Anisotropic transverse magnetoresistivity in α -YbAlB₄: Y. Matsumoto, J. Hong, K. Kuga and S. Nakatsuji, *J. Phys.: Conf. Ser.* **592** (2015) 012086(1-6).
15. Anomalous Enhancement of Seebeck Coefficient in Pr-Based 1-2-20 System with Non-Kramers Doublet Ground States: Y. Machida, T. Yoshida, T. Ikeura, K. Izawa, A. Nakama, R. Higashinaka, Y. Aoki, H. Sato, A. Sakai, S. Nakatsuji, N. Nagasawa, K. Matsumoto, T. Onimaru and T. Takabatake, *J. Phys.: Conf. Ser.* **592** (2015) 012025(1-9).
16. Anomalous specific heat behaviour in the quadrupolar Kondo system PrV₂Al₂₀: M. Tsujimoto, Y. Matsumoto and S. Nakatsuji, *J. Phys.: Conf. Ser.* **592** (2015) 012023.
17. *Antiferromagnetic transition of the caged compound TmTi₂Al₂₀: N. Kase, Y. Shimura, S. Kittaka, T. Sakakibara, S. Nakatsuji, T. Nakano, N. Takeda and J. Akimitsu, *J. Phys.: Conf. Ser.* **592** (2015) 012052(1-5).
18. *High Pressure Measurements of the Resistivity of β -YbAlB₄: T. Tomita, K. Kuga, Y. Uwatoko and S. Nakatsuji, *J. Phys.: Conf. Ser.* **592** (2015) 012019(1-5).
19. Shubnikov-de Haas Oscillation in the cubic Γ_3 -based heavy fermion superconductor PrV₂Al₂₀: Y. Shimura, M. Tsujimoto, A. Sakai, B. Zeng, L. Balicas and S. Nakatsuji, *J. Phys.: Conf. Ser.* **592** (2015) 012026(1-4).
20. *Synchrotron X-ray spectroscopy study on the valence state and magnetization in α -YbAl_{1-x}Fe_xB₄ ($x = 0.115$) at low temperatures and high magnetic fields: T. Terashima, Y. H. Matsuda, K. Kuga, S. Suzuki, Y. Matsumoto, S. Nakatsuji, A. Kondo, K. Kindo, N. Kawamura, M. Mizumaki and T. Inami, *J. Phys.: Conf. Ser.* **592** (2015) 012020 (6 pages).

† Joint research with outside partners.

21. ^{†*} Quadratic Fermi Node in a 3D Strongly Correlated Semimetal: T. Kondo, M. Nakayama, R. Chen, J. J. Ishikawa, E. -G. Moon, T. Yamamoto, Y. Ota, W. Malaeb, H. Kanai, Y. Nakashima, Y. Ishida, R. Yoshida, H. Yamamoto, M. Matsunami, S. Kimura, N. Inami, K. Ono, H. Kumigashira, S. Nakatsuji, L. Balents and S. Shin, *Nat. Commun.* **6** (2015) 10042 (1-8).
22. Foreword: Y. j. Kao, S. Onoda and S. Nakatsuji, *SPIN* **5** (2015) 1502001 (2 pages).
23. Special Issue on Quantum Spin Ice and Liquid in Geometrically Frustrated Magnets: Y. Kao, S. Onoda and S. Nakatsuji, *SPIN* **5** (2015) 1540001-1540006.
24. Unconventional Quantum Criticality in β -YbAlB₄ Detached from Its Magnetically Ordered Phase: T. Tomita, K. Kuga, Y. Uwatoko, P. Coleman and S. Nakatsuji, *Phys. Procedia* **75** (2015) 482-487.
25. ^{*} Field-induced quantum metal–insulator transition in the pyrochlore iridate Nd₂Ir₂O₇: Z. Tian, Y. Kohama, T. Tomita, H. Ishizuka, T. H. Hsieh, J. J. Ishikawa, K. Kindo, L. Balents and S. Nakatsuji, *Nature Phys.* **12** (2015) 134-139.
26. Pressure-Induced Local Structural Changes in Heavy Fermion β -YbAlB₄: Y. Sakaguchi, S. Ikeda, K. Kuga, S. Suzuki, S. Nakatsuji, N. Hirao, Y. Ohishi and H. Kobayashi, *J. Phys. Soc. Jpn.* **85** (2016) 023602(1-4).
27. Chemical and orbital fluctuations in Ba₃CuSb₂O₉: Y. Wakabayashi, D. Nakajima, Y. Ishiguro, K. Kimura, T. Kimura, S. Tsutsui, A. Q. R. Baron, K. Hayashi, N. Happo, S. Hosokawa, K. Ohwada and S. Nakatsuji, *Phys. Rev. B* **93** (2016) 245117(1-13).
28. Dimensional Reduction in Quantum Dipolar Antiferromagnets: P. Babkevich, M. Jeong, Y. Matsumoto, I. Kovacevic, A. Finco, R. Toft-Petersen, C. Ritter, M. Månsson, S. Nakatsuji and H. M. Rønnow, *Phys. Rev. Lett.* **116** (2016) 197202 (1-5).
29. ^{*} Experimental exploration of novel semimetal state in strong anisotropic Pyrochlore iridate Nd₂Ir₂O₇ under high magnetic field: Z. M. Tian, Y. Kohama, T. Tomita, J. Ishikawa, H. Mairo, K. Kindo and S. Nakatsuji, *J. Phys.: Conf. Ser.* **683** (2016) 012024(1-6).
30. ^{*} Frustrated magnetism in a Mott insulator based on a transition metal chalcogenide: S. Kawamoto, T. Higo, T. Tomita, S. Suzuki, Z. M. Tian, K. Mochizuki, A. Matsuo, K. Kindo and S. Nakatsuji, *J. Phys.: Conf. Ser.* **683** (2016) 012025 (1-4).
31. ^{*} Single crystal ²⁷Al-NMR study of the cubic Γ_3 ground doublet system PrTi₂Al₂₀: T. Taniguchi, M. Yoshida, H. Takeda, M. Takigawa, M. Tsujimoto, A. Sakai, Y. Matsumoto and S. Nakatsuji, *J. Phys.: Conf. Ser.* **683** (2016) 012016(1-9).
32. ^{*} Site-selective ¹¹B NMR studies on YbAlB₄: S. Takano, M. S. Grbic, K. Kimura, M. Yoshida, M. Takigawa, E. C. T. O. Farrell, K. Kuga, S. Nakatsuji and H. Harima, *J. Phys.: Conf. Ser.* **683** (2016) 012008(1-6).
33. Absence of Jahn-Teller transition in the hexagonal Ba₃CuSb₂O₉ single crystal: N. Katayama, K. Kimura, Y. Han, J. Nasu, N. Drichko, Y. Nakanishi, M. Halim, Y. Ishiguro, R. Satake, E. Nishibori, M. Yoshizawa, T. Nakano, Y. Nozue, Y. Wakabayashi, S. Ishihara, M. Hagiwara, H. Sawa and S. Nakatsuji, *Proc. Natl. Acad. Sci. U.S.A.* **112** (2015) 9305-9309.
34. Spin dynamics of the triangular-lattice antiferromagnet NiGa₂S₄ over an extended timescale: Y. Nambu, J. S. Gardner, D. E. MacLaughlin, C. Stock, H. Endo, S. Jonas, T. J. Sato, S. Nakatsuji and C. Broholm, *Phys. Rev. Lett.* **115** (2015) 127202(1-5).
35. ^{*} Low-Energy Excitations and Ground State Selection in Quantum Breathing Pyrochlore Antiferromagnet Ba₃Yb₂Zn₅O₁₁: T. Haku, K. Kimura, Y. Matsumoto, M. Soda, M. Sera, D. Yu, R. A. Mole, T. Takeuchi, S. Nakatsuji, Y. Kono, T. Sakakibara, L. -J. Chang and T. Masuda, *Phys. Rev. B* (2016), in print.

Division of Condensed Matter Theory

Takada group

Employing several standard techniques including the Green's-function method, quantum Monte Carlo simulations, band-structure calculations, and various types of variational approaches, we are studying several aspects of quantum many-body problems in condensed matter physics, based primarily on the first-principles Hamiltonian. This year we have studied the following issues: (1) The failure of the conventional self-consistent GW approximation in the calculation of both normal and superconducting properties is made clear by comparing the results among the G_0W_0 (one-shot GW), GW, and GWT approximations. (2) Diffusion Monte Carlo (DMC) simulations are performed on the system of an atom embedded in an electron gas with a view of investigating Kondo physics from first principles. A detailed analysis of the Friedel oscillations around the impurity atom

^{*} Joint research among groups within ISSP.

reveals that a proton-embedded electron gas can exhibit the Kondo temperature well beyond 1000 K. The obtained accurate electron-density profile is used to improve on the GGA-PBE version of the exchange-correlation energy functional in the density functional theory. In making this improvement, we have paid special attention to fulfilling the cusp theorem at the atom site. The improved functional will be applied to a wide range of topics in the future, including the phase diagram of the solid hydrogen under high pressures. (3) With proposing a better functional form for the vertex function Γ , always satisfying both the Ward identity and the momentum conservation law, we study the low-density electron gas in the GWT scheme to find an anomalous mass reduction as a result of avoiding the collapse of the normal state into a spontaneously excited electron-hole pair condensed state. Concomitantly with this mass reduction, an anomalous behavior of the momentum distribution function is found for the density parameter r_s around 20.

1. Emergence of a Kondo singlet state with Kondo temperature well beyond 1000 K in a proton-embedded electron gas: Y. Takada, R. Maezono and K. Yoshizawa, *Phys. Rev. B* **92** (2015) 155140(1-11).
2. Generic Features of an Electron Injected into the Luttinger Liquid: H. Maebashi and Y. Takada, *J. Supercond. Nov. Magn.* **28** (2015) 1331-1335.
3. Role of the ward identity and relevance of the G^0W^0 approximation in normal and superconducting states: Y. Takada, *Mol. Phys.* **114** (2016) 1.
4. Theory for Reliable First-Principles Prediction of the Superconducting Transition Temperature: Y. Takada, in: *Carbon-based New Superconductors: Toward high-Tc superconductivity (ISBN 978-981-4303-30-9 (Hardcover), 978-981-4303-31-6 (eBook))*, Ch 8, edited by J. Haruyama, (Pan Stanford Publishing Pte. Ltd., 2015), pp. 193-230.
5. Reference Module in Materials Science and Materials Engineering: Y. Takada, in: *Materials Science and Materials Engineering*, edited by S. Mahfoudh and M. Nicholls, (Elsevier, 2016), B9780128035818007748.

Oshikawa group

We studied a wide range of fundamental problems in condensed matter theory and statistical mechanics. In particular, we discovered that even trivial phases, which are adiabatically connected to a product state without any entanglement, are not unique in the presence of an appropriate symmetry. We discussed a simple example of $S = 1$ quantum spin chain with a symmetry under a combined operation of the site-centered lattice inversion and the global π -rotation about z axis. In this model, there are two trivial phases, adiabatically connected to the Néel state and the large-D state (product of $S^z=0$ states), which are always separated by a quantum phase transition in the presence of the above symmetry. We demonstrated this using field theory, numerical calculation, and a general proof based on Matrix Product State representation. The present result brings about a new perspective in classification of quantum phases, a central issue in current condensed matter physics.

1. Equilibrium surface current and role of $U(1)$ symmetry: Sum rule and surface perturbations: Y. Tada, *Phys. Rev. B* **92** (2015) 104502(1-15).
2. [†]Selection of factorizable ground state in a frustrated spin tube: Order by disorder and hidden ferromagnetism: X. Plat, Y. Fuji, S. Capponi and P. Pujol, *Phys. Rev. B* **91** (2015) 064411 (1-21).
3. [†]Spin fluctuations and superconductivity in layered f-electron superlattices: Y. Tada and R. Peters, *Phys. Rev. B* **92** (2015) 035129(1-8).
4. [†]Absence of Quantum Time Crystals: H. Watanabe and M. Oshikawa, *Phys. Rev. Lett.* **114** (2015) 251603(1-5).
5. [†]Distinct Trivial Phases Protected by a Point-Group Symmetry in Quantum Spin Chains: Y. Fuji, F. Pollmann and M. Oshikawa, *Phys. Rev. Lett.* **114** (2015) 177204.
6. Gravitational Casimir Effect: J. Q. Quach, *Phys. Rev. Lett.* **114** (2015) 081104(1-5).
7. [†]Orbital Angular Momentum and Spectral Flow in Two-Dimensional Chiral Superfluids: Y. Tada, W. Nie and M. Oshikawa, *Phys. Rev. Lett.* **114** (2015) 195301.
8. 「物性物理学」～物質に潜む普遍の概念～: 押川 正毅, *数理科学* **620** (2015) 40-45.
9. Foldy-Wouthuysen transformation of the generalised Dirac Hamiltonian in a gravitational-wave background: J. Q. Quach, *Physical Review D* **92** (2015) 084047.
10. [†]Flux quench in a system of interacting spinless fermions in one dimension: Y. O. Nakagawa, G. Misguich and M. Oshikawa, *Phys. Rev. B* **93** (2016) 174310.
11. [†]Magnetism and superconductivity in ferromagnetic heavy-fermion system UCoGe under in-plane magnetic fields: Y. Tada, S. Takayoshi and S. Fujimoto, *Phys. Rev. B* **93** (2016) 174512(1-7).

[†] Joint research with outside partners.

12. [†]Plaquette order in the SU(6) Heisenberg model on the honeycomb lattice: P. Nataf, M. Lajkó, P. Corboz, A. M. Läuchli, K. Penc and F. Mila, Phys. Rev. B **93** (2016) 201113(R)(1-6).
13. Spin gravitational resonance and graviton detection: J. Q. Quach, Physical Review D **93** (2016) 104048(1-6).
14. [†]ウラン系強磁性超伝導における強磁性ゆらぎが誘起するスピン三重項超伝導: 石田 憲二, 服部 泰佑, 佐藤 憲昭, 出口 和彦, 多田 靖啓, 藤本 聡, 固体物理 **50** (2015) 123-132.
15. 量子異常と物性物理 (特集 物理科学, この1年) - (素粒子物理): 押川 正毅, パリティ **31** (2016) 43-45.
16. 物性物理と場の量子論 ～場の量子論が明かす双対性～: 押川 正毅, 数理科学 **633** (2016) 41-46.

Tsunetsugu group

We have studied optical conductivity near an antiferromagnetic phase transition in the square-lattice Hubbard model at half filling using the cluster dynamical-mean field approach. We investigated the effects of vertex corrections on optical conductivity and found that they have large contributions and change some important features of optical conductivity. The vertex corrections enhance frequency dependence of conductivity in both metallic and insulating phases. Another important discovery is the presence of a temperature region above the transition temperature where dc conductivity shows non-increasing behavior with lowering temperature. This is not due to a pseudogap behavior, but the electron spectral function does not show a dip at Fermi energy. (Reference: arXiv:1605.00387) We have continued the study of quadrupole order in Pr 1-2-20 system. We used a classical Monte Carlo calculation to investigate the effects of thermal fluctuations in temperature-magnetic field phase diagram. We found that the same number of ordered phases appear as predicted by our previous mean-field analysis, but the phase boundaries are strongly modified by thermal fluctuations and have a different topology. A new tetracritical point appears when magnetic field is applied along (001) direction. Criticality of parasitic ferro quadrupole order is also investigated, and we have found an unusual critical behavior in their temperature dependence. (Reference arXiv:1605.05175) We have also studied an antiferromagnetic Heisenberg model on breathing pyrochlore lattice, and found that the ground state in the $S = 3/2$ case has an interesting "dimerized" pattern that differs from the previously studied $S = 1/2$ case.

1. Fano resonance through Higgs bound states in tunneling of Nambu-Goldstone modes: T. Nakayama, I. Danshita, T. Nikuni and S. Tsuchiya, Phys. Rev. A **92** (2015) 043610(1-19).
2. Dynamical Characteristics of the Mott Transition: Examination of Doublon Dynamics in a Triangular-lattice Hubbard Model: T. Sato and H. Tsunetsugu, Physics Procedia **75** (2015) 376-382.

Sugino group

We have done first-principles study of the excited-states, ferroelectric interfaces, electrochemical interfaces, and the ground-state wavefunctions. We have improved our GW+ BSE program of excited-state calculation so as to manipulate large number of atoms (~200 atoms), which has significantly activated collaboration with experiments. We also advanced density functional methods to compute the electrically biased interface, which are used to investigate the system of negative capacitance as well as the electrochemical reactions. We further developed a variational approach to strongly correlated electron systems to investigate the structure of the many-body wavefunction.

1. All-electron GW+Bethe-Salpeter calculations on small molecules: D. Hirose, Y. Noguchi and O. Sugino, Phys. Rev. B **91** (2015) 205111.
2. [†]First-principles calculation of charged capacitors under open-circuit conditions using the orbital-separation approach: S. Kasamatsu, S. Watanabe and S. Han, Phys. Rev. B **92** (2015) 115124.
3. Symmetry breaking and excitonic effects on optical properties of defective nanographenes: Y. Noguchi and O. Sugino, J. Chem. Phys. **142** (2015) 064313.
4. Configuration interaction with antisymmetrized geminal powers: W. Uemura, S. Kasamatsu and O. Sugino, Phys. Rev. A **91** (2015) 062504.
5. *Vibronic Structures in Absorption and Fluorescence Spectra of Firefly Oxyluciferin in Aqueous Solutions: M. Hiyama, Y. Noguchi, H. Akiyama, K. Yamada and N. Koga, Photochem. Photobiol. **91** (2015) 819.
6. *First-Principles Investigation of Strong Excitonic Effects in Oxygen 1s X-ray Absorption Spectra: Y. Noguchi, M. Hiyama, H. Akiyama, Y. Harada and N. Koga, J. Chem. Theory Comput. **11** (2015) 1668-1673.
7. [†]TOMBO: All-electron mixed-basis approach to condensed matter physics: S. Ono, Y. Noguchi, R. Sahara, Y. Kawazoe and K. Ohno, Comput. Phys. Commun. **189** (2015) 20.

* Joint research among groups within ISSP.

8. ^{*} Pressure dependence of the magnetic ground states in MnP: M. Matsuda, F. Ye, S. E. Dissanayake, J. -G. Cheng, S. Chi, J. Ma, H. D. Zhou, J. -Q. Yan, S. Kasamatsu, O. Sugino, T. Kato, K. Matsubayashi, T. Okada and Y. Uwatoko, *Phys. Rev. B* **93** (2016) 100405(1-5).
9. [†] Emergence of Negative Capacitance in Multidomain Ferroelectric-Paraelectric Nanocapacitors at Finite Bias: S. Kasamatsu, S. Watanabe, C. S. Hwang and S. Han, *Adv. Mater.* **28** (2016) 335.

Kato group

The main research subject of our laboratory is theory of non-equilibrium properties in nanoscale devices. We have studied (1) photon-assisted current noises under strong AC fields in quantum dot systems, (2) a multi-orbital Anderson impurity at high bias voltages, and (3) non-equilibrium current noises of quantum dots in the Kondo regime. We have also studied form factors of the Kondo model by the Bethe ansatz method.

1. Effects of Coulomb interaction on photon-assisted current noise through a quantum dot: T. J. Suzuki and T. Kato, *Phys. Rev. B* **91** (2015) 165302(1-12).
2. Exact Green's function for a multiorbital Anderson impurity at high bias voltages: A. Oguri and R. Sakano, *Phys. Rev. B* **91** (2015) 115429(1-14).
3. Universality of non-equilibrium fluctuations in strongly correlated quantum liquids: M. Ferrier, T. Arakawa, T. Hata, R. Fujiwara, R. Delagrè, R. Weil, R. Deblock, R. Sakano, A. Oguri and K. Kobayashi, *Nature Phys.* **12** (2015) 230-235.

Division of Nanoscale Science

Iye group

Electronic transport in monolayer graphene grown on vicinal surface of 6H-SiC(0001) with a quasi-regular step-and-terrace structure is studied. Conductivity under a magnetic field normal to the plane showed a high degree of anisotropy. The quantum Hall effect (QHE) with zero resistance manifests itself for the current along the steps, whereas the QHE is obscured by pronounced positive magnetoresistance with quadratic magnetic-field dependence for the current across the steps.

1. [†] Ballistic transport in graphene antidot lattices: R. Yagi, R. Sakakibara, R. Ebisuoka, J. Onishi, K. Watanabe, T. Taniguchi and Y. Iye, *Phys. Rev. B* **92** (2015) 195406(1-6).
2. ^{†*} Highly Anisotropic Parallel Conduction in the Stepped Substrate of Epitaxial Graphene Grown on Vicinal SiC: A. Endo, F. Komori, K. Morita, T. Kajiwarra and S. Tanaka, *J. Low Temp. Phys.* **179** (2015) 237-250.

Katsumoto group

Conductance fluctuation in InAs two-dimensional electrons versus in-plane magnetic field was found and attributed to the sign of so called "Zitterbewegung" (trembling motion) due to spin-orbit coupling. We have succeeded in making low resistance semiconductor-superconductor junctions not only for InAs two-dimensional electrons but also for a diluted magnetic semiconductor (In,Fe)As. Anomalous response in the latter will be our next subject.

1. [†] Gate-Tunable Atomically Thin Lateral MoS₂ Schottky Junction Patterned by Electron Beam: Y. Katagiri, T. Nakamura, A. Ishii, C. Ohata, M. Hasegawa, S. Katsumoto, T. Cusati, A. Fortunelli, G. Iannaccone, G. Fiori, S. Roche and J. Haruyama, *Nano Lett.* **16** (2016) 3788-3794.

Otani group

We have studied on three topics including spin Hall effect, spin diffusion length, and magnonic crystals.

1. Tunable configurational anisotropy in collective magnetization dynamics of Ni₈₀Fe₂₀ nanodot arrays with varying dot shapes: B. K. Mahato, S. Choudhury, R. Mandal, S. Barman, Y. Otani and A. Barman, *J. Appl. Phys.* **117** (2015) 213909.
2. Tunable spin wave spectra in two-dimensional Ni₈₀Fe₂₀ antidot lattices with varying lattice symmetry: R. Mandal, S. Barman, S. Saha, Y. Otani and A. Barman, *J. Appl. Phys.* **118** (2015) 053910.
3. Crossover between spin swapping and Hall effect in disordered systems: H. B. M. Saidaoui, Y. Otani and A. Manchon, *Phys. Rev. B* **92** (2015) 024417.

[†] Joint research with outside partners.

4. Revisiting the measurement of the spin relaxation time in graphene-based devices: H. Idzuchi, A. Fert and Y. Otani, *Phys. Rev. B* **91** (2015) 241407.
5. Spin relaxation mechanism in a highly doped organic polymer film: M. Kimata, D. Nozaki, Y. Niimi, H. Tajima and Y. Otani, *Phys. Rev. B* **91** (2015) 224422.
6. Strong Suppression of the Spin Hall Effect in the Spin Glass State: Y. Niimi, M. Kimata, Y. Omori, B. Gu, T. Ziman, S. Maekawa, A. Fert and Y. Otani, *Phys. Rev. Lett.* **115** (2015) 196602.
7. スピン軌道相互作用の強い導体中のスピホール効果: 大谷 義近, 新見 康洋, *固体物理* **50** (2015) 575-590.
8. Spin relaxation characteristics in Ag nanowire covered with various oxides: S. Karube, H. Idzuchi, K. Kondou, Y. Fukuma and Y. Otani, *Appl. Phys. Lett.* **107** (2015) 122406.
9. Selective mode excitation in three-chained magnetic vortices: N. Hasegawa, S. Sugimoto, H. Fujimori, K. Kondou, Y. Niimi and Y. Otani, *Appl. Phys. Express* **8** (2015) 063005-1.
10. Reciprocal spin Hall effects in conductors with strong spin-orbit coupling: a review: Y. Niimi and Y. Otani, *Rep. Prog. Phys.* **78** (2015) 124501.
11. Spin transport in non-magnetic nano-structures induced by non-local spin injection: H. Idzuchi, Y. Fukuma and Y. Otani, *Physica E: Low-dimensional Systems and Nanostructures* **68** (2015) 239.
12. Realization of a micrometre-scale spin-wave interferometer: O. Rousseau, B. Rana, R. Anami, M. Yamada, K. Miura, S. Ogawa and Y. Otani, *Sci. Rep.* **5** (2015) 9873.
13. Quasiparticle-mediated spin Hall effect in a superconductor: T. Wakamura, H. Akaike, Y. Omori, Y. Niimi, S. Takahashi, A. Fujimaki, S. Maekawa and Y. Otani, *Nature Mater.* **14** (2015) 675.
14. Tunable picosecond spin dynamics in two dimensional ferromagnetic nanodot arrays with varying lattice symmetry: S. Saha, S. Barman, S. Sugimoto, Y. Otani and A. Barman, *RSC Adv.* **5** (2015) 34027.
15. All-optical investigation of tunable picosecond magnetization dynamics in ferromagnetic nanostripes with a width down to 50 nm: S. Saha, S. Barman, Y. Otani and A. Barman, *Nanoscale* **7** (2015) 18312.

Komori group

Reproducible dependence of the STM images on the distance between the surface and the STM tip apex for a monatomic layer of iron nitride (Fe_2N) formed on a Cu(001) surface was found in the bias-voltage range corresponding to the Fe $3d$ states. The results are attributed to a shift in surface orbitals detected by the tip from the d states to the sp states. Electronic structures of the Pt-adsorbed Ge(001) surface with a one-dimensional atomic structure were studied by high-resolution ARPES and SARPES. One-dimensional Fermi surfaces of four surface states were confirmed by ARPES. One of them is spin-split due to the Rashba effect.

1. [†]Nonlinear terahertz field-induced carrier dynamics in photoexcited epitaxial monolayer graphene: H. A. Hafez, I. Al-Naib, M. M. Dignam, Y. Sekine, K. Oguri, F. Blanchard, D. G. Cooke, S. Tanaka, F. Komori, H. Hibino and T. Ozaki, *Phys. Rev. B* **91** (2015) 035422 (1-9).
2. ^{*}Scanning tunneling spectroscopy study of quasiparticle interference on dual topological insulator $\text{Bi}_{1-x}\text{Sb}_x$: S. Yoshizawa, F. Nakamura, A. A. Taskin, T. Iimori, K. Nakatsuji, I. Matsuda, Y. Ando and F. Komori, *Phys. Rev. B* **91** (2015) 045423(1-6).
3. ^{†*}Highly Anisotropic Parallel Conduction in the Stepped Substrate of Epitaxial Graphene Grown on Vicinal SiC: A. Endo, F. Komori, K. Morita, T. Kajiwara and S. Tanaka, *J. Low Temp. Phys.* **179** (2015) 237-250.
4. ^{†*}フェムト秒域時間分解光電子分光法によるグラフェンの超高速キャリアダイナミクスの追跡 Tracing Ultrafast Carrier Dynamics in Graphene with Femtosecond Time-resolved Photoemission Spectroscopy: 染谷 隆史, 吹留 博一, 石田 行章, 吉田 力矢, 山本 達, 板谷 治郎, 小森 文夫, 辛 埴, 松田 巖, *表面科学* **36(8)** (2015) 418-423.
5. ^{†*}Layer number dependence of carrier lifetime in graphenes observed using time-resolved mid-infrared luminescence: H. Watanabe, T. Kawasaki, T. Iimori, F. Komori and T. Suemoto, *Chem. Phys. Lett.* **637** (2015) 58-62.
6. ^{†*}Selective Formation of Zigzag Edges in Graphene Cracks: M. Fujihara, R. Inoue, R. Kurita, T. Taniuchi, Y. Motoyui, S. Shin, F. Komori, Y. Maniwa, H. Shinohara and Y. Miyata, *ACS Nano* **9** (2015) 9027-9033.

^{*} Joint research among groups within ISSP.

7. ^{†*}One-dimensional metallic surface states of Pt-induced atomic nanowires on Ge(001): K. Yaji, S. Kim, I. Mochizuki, Y. Takeichi, Y. Ohtsubo, P. L. Fèvre, F. Bertran, A. Taleb-Ibrahimi, S. Shin and F. Komori, *J. Phys.: Condens. Matter* **28** (2016) 284001(1-9).
8. *High-resolution three-dimensional spin- and angle-resolved photoelectron spectrometer using vacuum ultraviolet laser light: K. Yaji, A. Harasawa, K. Kuroda, S. Toyohisa, M. Nakayama, Y. Ishida, A. Fukushima, S. Watanabe, C. Chen, F. Komori and S. Shin, *Rev. Sci. Instrum.* **87** (2016) 053111(1-6).
9. Orbital Selectivity in Scanning Tunneling Microscopy: Distance-Dependent Tunneling Process Observed in Iron Nitride: Y. Takahashi, T. Miyamachi, K. Ienaga, N. Kawamura, A. Ernst and F. Komori, *Phys. Rev. Lett.* **116** (2016) 056802(1-5).
10. ^{†*}Spin Polarization and Texture of the Fermi Arcs in the Weyl Fermion Semimetal TaAs: S.-Y. Xu, I. Belopolski, D. S. Sanchez, M. Neupane, G. Chang, K. Yaji, Z. Yuan, C. Zhang, K. Kuroda, G. Bian, C. Guo, H. Lu, T.-R. Chang, N. Alidoust, H. Zheng, C.-C. Lee, S.-M. Huang, C.-H. Hsu, H.-T. Jeng, A. Bansil, T. Neupert, F. Komori, T. Kondo, S. Shin, H. Lin, S. Jia and M. Zahid Hasan, *Phys. Rev. Lett.* **116** (2016) 096801(1-7).
11. Ribbon-Like Nanopattern Formed on Nitrogen-Adsorbed Vicinal Cu(001): M. Yamada, N. Kawamura, K. Nakatsuji and F. Komori, *e-J. Surf. Sci. Nanotech.* **14** (2016) 43-46.
12. *Photoelectrochemical water splitting enhanced by self-assembled metal nanopillars embedded in an oxide semiconductor photoelectrode: S. Kawasaki, R. Takahashi, T. Yamamoto, M. Kobayashi, H. Kumigashira, J. Yoshinobu, F. Komori, A. Kudo and M. Lippmaa, *Nat. Commun.* **7** (2016) 11818(1-6).

Yoshinobu group

We conducted several research projects in the fiscal year 2015. (1) The activation and hydrogenation of CO₂ on clean and Zn-deposited Cu(111) and Cu(997) surfaces studied by AP-XPS. (2) The surface chemistry of formic acid on Zn-Cu(111) studied by SR-PES. (3) Spectroscopic characterization of H-Cu(111), Zn-Cu(111) and Pd-Cu surfaces by XPS. (4) Spectroscopic characterization of Au on SrTiO₃ under O₂ exposure using SR-XPS (5) LT-STM study of CO₂ on Cu(997) (6) Independently driven four-probe conductivity measurement of organic thin films.

1. Quantitative analysis of desorption and decomposition kinetics of formic acid on Cu(111): the importance of hydrogen bonding between adsorbed species: Y. Shiozawa, T. Koitaya, K. Mukai, S. Yoshimoto and J. Yoshinobu, *J. Chem. Phys.* **143** (2015) 234707.
2. Observation of Fano line shapes in infrared vibrational spectra of CO₂ adsorbed on Cu(997): T. Koitaya, Y. Shiozawa, K. Mukai, S. Yoshimoto and J. Yoshinobu, *J. Chem. Phys.* **144** (2016) 054703.
3. *Photoelectrochemical water splitting enhanced by self-assembled metal nanopillars embedded in an oxide semiconductor photoelectrode: S. Kawasaki, R. Takahashi, T. Yamamoto, M. Kobayashi, H. Kumigashira, J. Yoshinobu, F. Komori, A. Kudo and M. Lippmaa, *Nat. Commun.* **7** (2016) 11818(1-6).
4. *Real-time observation of reaction processes of CO₂ on Cu(997) by ambient-pressure X-ray photoelectron spectroscopy: T. Koitaya, S. Yamamoto, Y. Shiozawa, K. Takeuchi, R.-Y. Liu, K. Mukai, S. Yoshimoto, K. Akikubo, I. Matsuda and J. Yoshinobu, *Topic in Catalysis* **59** (2016) 526-531.
5. The chemistry of simple alkene molecules on Si(100)c(4x2): the mechanism of cycloaddition and their selectivities: K. Akagi and J. Yoshinobu, *Surface Science* (2016), accepted for publication.

Hasegawa group

We have developed spin-polarized scanning tunneling microscopy (SP-STM), which enables us to detect the orientation of spins / magnetic moments on surfaces in atomic-scale spatial resolutions. Using the method, we investigated the formation mechanism of spin spiral structures of Mn ultrathin films formed on a W(110) substrate. Because of the absence of inversion symmetry due to the thin film structure and the spin-orbit interaction in the heavy-elemental substrate, the Dzyaloshinskii-Moriya interaction (DMI) is exerted among the spins in the thin films, which induces chirality in the spin structures. In order to investigate the details of DMI, we directly measured the rotational sense of the spin spiral structures by SP-STM, and revealed that both 1st and 2nd Mn layers exhibit chirality and the same polarization of DMI despite their different types of spin structures and propagation directions. These results, combined with previous reports on the chirality of domain walls observed in magnetic thin films on the same substrate, lead us to conclude that the DMI polarization is dominantly determined by the substrate. Aiming for a bottom-up approach of fabricating quantum spin systems and their microscopic investigations, we studied the adsorption of oxygen molecules, which have a spin triplet state ($S = 1$) as a ground state, on a Ag(111) substrate. It was found that by low temperature exposure the molecules adsorb on the substrate in a physisorbed manner lying down to form a triangular

[†] Joint research with outside partners.

lattice. Different from an isosceles triangular lattice expected from the ellipsoidal shape of the molecule, the lattice is deformed to scalene. Based on a Monte Carlo calculation using parameters that account for the solid oxygen phases, we found that the introduction of antiferromagnetic interaction among the adsorbed molecules explain the deformation quantitatively. The antiferromagnetic order indicates the preservation of the spin of the adsorbed molecules, opening up the possibility that the system can be utilized for fabrication / construction of one- or two- dimensional spin structures using an atom / molecular manipulation method of STM.

1. Disorder-induced suppression of superconductivity in the Si(111)-($\sqrt{7} \times \sqrt{3}$)-In surface: Scanning tunneling microscopy study: S. Yoshizawa, H. Kim, Y. Hasegawa and T. Uchihashi, *Phys. Rev. B* **92** (2015) 041410(R)(1-5).
2. Electronic and magnetic effects of a stacking fault in cobalt nanoscale islands on the Ag(111) surface: K. Doi, E. Minamitani, S. Yamamoto, R. Arafune, Y. Yoshida, S. Watanabe and Y. Hasegawa, *Phys. Rev. B* **92** (2015) 064421(1-8).
3. Site-Dependent Evolution of Electrical Conductance from Tunneling to Atomic Point Contact: H. Kim and Y. Hasegawa, *Phys. Rev. Lett.* **114** (2015) 206801(1-5).
4. バルク Cr 探針を用いた W(110) 上の Mn 薄膜のスピン偏極 STM/STS 観察: 土師 将裕, 吉田 靖雄, 長谷川 幸雄, *表面科学* **36** (2015) 403-407.
5. [†]Impact of Surface Conditions on the Superconductivity of Si(111)-($\sqrt{7} \times \sqrt{3}$)-In: S. Yoshizawa, H. Kim, T. Kawakami, Y. Nagai, T. Nakayama, X. Hu, Y. Hasegawa and T. Uchihashi, *e-J. Surf. Sci. Nanotech.* **13** (2015) 151-154.
6. Direct visualization of surface phase of oxygen molecules physisorbed on Ag(111) surface: A two-dimensional quantum spin system: S. Yamamoto, Y. Yoshida, H. Imada, Y. Kim and Y. Hasegawa, *Phys. Rev. B* **93** (2016) 081408(R)(1-5).
7. Insensitivity of atomic point contact conductance to a moiré structure: H. Kim and Y. Hasegawa, *Phys. Rev. B* **93** (2016) 075409(1-6).
8. Spatial variation in local work function as an origin of moiré contrast in scanning tunneling microscopy images of Pb thin films / Si(111): H. Kim and Y. Hasegawa, *Jpn. J. Appl. Phys.* (2016), accepted for publication.
9. Superconducting proximity effect on Rashba-split Pb/Ge(111)- $\beta\sqrt{3} \times \sqrt{3}$ surface: H. Kim, Y. Miyata and Y. Hasegawa, *Supercond. Sci. Technol.* (2016), accepted for publication.

Lippmaa group

We work on various aspects of oxide thin films and interfaces. One of our aims is to explore novel polar oxide phases that may exhibit new type of magnetoelectric coupling. In particular, we have looked at double perovskites that can sustain both ferromagnetism and ferroelectricity in a strained lattice. We have continued working on magnetic interfaces, mostly looking at band alignment and magnetic coupling in manganite-titanate heterostructures. The third topic is the development of photocatalytic materials. Our recent work has involved the analysis of carrier dynamics in optically excited crystals, surface chemical reactions, and the development of nanoscale composite materials for efficient collection of photogenerated charge in a photocatalyst.

1. Interfacial capacitance between a ferroelectric Fe₃O₄ thin film and a semiconducting Nb:SrTiO₃ substrate: R. Takahashi, Y. Cho and M. Lippmaa, *J. Appl. Phys.* **117** (2015) 014104(1-10).
2. A-site-driven ferroelectricity in strained ferromagnetic La₂NiMnO₆ thin films: R. Takahashi, I. Ohkubo, K. Yamauchi, M. Kitamura, Y. Sakurai, M. Oshima, T. Oguchi, Y. Cho and M. Lippmaa, *Phys. Rev. B* **91** (2015) 134107(1-9).
3. [†]Determination of band diagram for a p-n junction between Mott insulator LaMnO₃ and band insulator Nb:SrTiO₃: M. Kitamura, M. Kobayashi, E. Sakai, R. Takahashi, M. Lippmaa, K. Horiba, H. Fujioka and H. Kumigashira, *Appl. Phys. Lett.* **106** (2015) 061605(1-5).
4. Magnetic coupling at perovskite and rock-salt structured interfaces: M. Matvejeff, E. Ahvenniemi, R. Takahashi and M. Lippmaa, *Appl. Phys. Lett.* **107** (2015) 141604(1-5).
5. [†]Photo-electrochemical epitaxy of silver-oxide clathrate Ag₇O₈M (M=NO₃, HSO₄) on rutile-type Nb-doped TiO₂ single crystals: R. Tanaka, R. Takahashi, S. Takata, M. Lippmaa and Y. Matsumoto, *CrystEngComm* **17** (2015) 3701-3707.
6. *Optical pump-THz probe analysis of long-lived *d*-electrons and relaxation to self-trapped exciton states in MnO: J. Nishitani, T. Nagashima, M. Lippmaa and T. Suemoto, *Appl. Phys. Lett.* **108** (2016) 162101(1-5).
7. *Photoexcited *d*-electron dynamics in transition metal oxide MnO studied by optical pump-THz probe measurements: J. Nishitani, T. Kurihara, A. Asahara, T. Nagashima, M. Lippmaa and T. Suemoto, *Phys. Status Solidi C* **13** (2016) 113-116.

* Joint research among groups within ISSP.

8. The effect of polar (111)-oriented SrTiO₃ on initial perovskite growth: I. Hallsteinsen, M. Nord, T. Bolstad, P. -E. Vullum, J. E. Boschker, P. Longo, R. Takahashi, R. Holmestad, M. Lippmaa and T. Tybell, *Cryst. Growth Des.* **16** (2016) 2357-2362.
9. *Photoelectrochemical water splitting enhanced by self-assembled metal nanopillars embedded in an oxide semiconductor photoelectrode: S. Kawasaki, R. Takahashi, T. Yamamoto, M. Kobayashi, H. Kumigashira, J. Yoshinobu, F. Komori, A. Kudo and M. Lippmaa, *Nat. Commun.* **7** (2016) 11818(1-6).

Division of Physics in Extreme Conditions

Uwatoko group

We report electrical resistivity, ac magnetic susceptibility and X-ray absorption spectroscopy measurements of intermediate valence YbNi₃Ga₉ under pressure and magnetic field. We have revealed a characteristic pressure-induced Yb valence crossover within the temperature-pressure phase diagram, and a first-order metamagnetic transition is found below $P_c \sim 9$ GPa where the system undergoes a pressure-induced antiferromagnetic transition. Zirconium-based bulk metallic glass (Zr-based BMG) has outstanding properties as a cylinder material for piston-cylinder high pressure apparatuses and is especially useful for neutron scattering. The piston cylinder consisting of a Zr-based BMG cylinder with outer/inner diameters of 8.8/2.5 mm sustains pressures up to 1.81 GPa and ruptured at 2.0 GPa, with pressure values determined by the superconducting temperature of lead. We report the discovery of pressure-induced superconductivity below $T_c = 14$ K in the iron-based spin-ladder material BaFe₂S₃, a Mott insulator with striped-type magnetic ordering below ~ 120 K. Our findings indicate that iron-based ladder compounds represent promising material platforms, in particular for studying the fundamentals of iron-based superconductivity. The perovskite PbCrO₃ is an antiferromagnetic insulator. However, the fundamental interactions leading to the insulating state in this single valent perovskite are unclear. We report a variety of insitu pressure measurements including electron transport properties, X-ray absorption spectrum, and crystal structure study by X-ray and neutron diffraction. These studies reveal key information leading to the elucidation of the physics behind the insulating state and the pressure-induced transition.

1. †Electron transport in TTF-CA under High pressures: R. Takehara, K. Miyagawa, K. Kanoda, T. Miyamoto, H. Matsuzaki, H. Okamoto, H. Taniguchi, K. Matsubayashi and Y. Uwatoko, *Physica B* **460** (2015) 83-87.
2. †* Anomalous Quantum Transport Properties in Semimetallic Black Phosphorus: K. Akiba, A. Miyake, Y. Akahama, K. Matsubayashi, Y. Uwatoko, H. Arai, Y. Fuseya and M. Tokunaga, *J. Phys. Soc. Jpn.* **84** (2015) 073708(1-4).
3. †Correlation between T_c and Crystal Structure in S-Doped FeSe Superconductors under Pressure: Studied by X-ray Diffraction of FeSe_{0.8}S_{0.2} at Low Temperatures: T. Tomita, H. Takahashi, H. Okada, Y. Mizuguchi, Y. Takano, S. Nakano, K. Matsubayashi and Y. Uwatoko, *J. Phys. Soc. Jpn.* **84** (2015) 024713(1-8).
4. †Electrical Transport in the Quasi-Two-Dimensional Ionic Mott Insulator M₂P-TCNQF₄ under High Pressures: R. Takehara, K. Miyagawa, T. Miyamoto, H. Okamoto, H. Taniguchi, K. Matsubayashi, Y. Uwatoko and K. Kanoda, *J. Phys. Soc. Jpn.* **84** (2015) 104702(1-5).
5. †High Pressure Effect on the Superconductivity in VN: B. Wang, K. Matsubayashi, Y. Uwatoko and K. Ohgushi, *J. Phys. Soc. Jpn.* **84** (2015) 104706(1-4).
6. †*Magnetic Order in the Frustrated Ising Quasi-One Dimensional Compound NaCo(acac)₃·Benzene: Y. Karaki, K. Kuga, K. Kimura, S. Nakatsuji, K. Matsubayashi and Y. Uwatoko, *J. Phys. Soc. Jpn.* **84** (2015) 084708.
7. †Pressure-Induced Valence Transition and Heavy Fermion State in Eu₂Ni₃Ge₅ and EuRhSi₃: A. Nakamura, T. Okazaki, M. Nakashima, Y. Amako, K. Matsubayashi, Y. Uwatoko, S. Kayama, T. Kagayama, K. Shimizu, T. Uejo, H. Akamine, M. Hedo, T. Nakama, Y. Onuki and H. Shiba, *J. Phys. Soc. Jpn.* **84** (2015) 053701(1-4).
8. †Transport and Magnetic Properties of EuAl₄ and EuGa₄: A. Nakamura, T. Uejo, F. Honda, T. Takeuchi, H. Harima, E. Yamamoto, Y. Haga, K. Matsubayashi, Y. Uwatoko, M. Hedo, T. Nakama and Y. Onuki, *J. Phys. Soc. Jpn.* **84** (2015) 124711(1-9).
9. Upper Critical Field, Critical Current Density and Activation Energy of the New La_{1-x}Sm_xO_{0.5}F_{0.5}BiS₂(x= 0.2, 0.8) Superconductors: G. K. Selvan, G. S. Thakur, K. Manikandan, Y. Uwatoko, Z. Haque, L. C. Gupta, A. K. Ganguli and S. Arumugam, *J. Phys. Soc. Jpn.* **84** (2015) 124701(1-5).
10. Ferromagnetic superexchange in insulating Cr₂MoO₆ by controlling orbital hybridization: M. Zhu, D. Do, C. R. Dela Cruz, Z. Dun, J. -G. Cheng, H. Goto, Y. Uwatoko, T. Zou, H. D. Zhou, S. D. Mahanti and X. Ke, *Phys. Rev. B* **92** (2015) 094419(1-6).

† Joint research with outside partners.

11. * Temperature and composition phase diagram in the iron-based ladder compounds $\text{Ba}_{1-x}\text{Cs}_x\text{Fe}_2\text{Se}_3$: T. Hawaii, Y. Nambu, K. Ohgushi, F. Du, Y. Hirata, M. Avdeev, Y. Uwatoko, Y. Sekine, H. Fukazawa, J. Ma, S. Chi, Y. Ueda, H. Yoshizawa and T. J. Sato, *Phys. Rev. B* **91** (2015) 184416.
12. Pressure Induced Superconductivity on the border of Magnetic Order in MnP: J. -G. Cheng, K. Matsubayashi, W. Wu, J. P. Sun, F. K. Lin, J. L. Luo and Y. Uwatoko, *Phys. Rev. Lett.* **114** (2015) 117001(1-18).
13. Pressure-Induced Valence Crossover and Novel Metamagnetic Behavior near the Antiferromagnetic Quantum Phase Transition of YbNi_3Ga_9 : K. Matsubayashi, T. Hirayama, T. Yamashita, S. Ohara, N. Kawamura, M. Mizumaki, N. Ishimatsu, S. Watanabe, K. Kitagawa and Y. Uwatoko, *Phys. Rev. Lett.* **114** (2015) 086401(1-5).
14. Development of High-Pressure and Multi-Frequency ESR System and Its Application to Quantum Spin System: T. Sakurai, R. Matsui, K. Kawasaki, S. Okubo, H. Ohta, K. Matsubayashi, Y. Uwatoko, K. Kudo and Y. Koike, *Appl. Magn. Reson.* **46** (2015) 1007-1012.
15. †Development of High-Pressure ESR System Using Micro-coil: K. Kawasaki, T. Sakurai, E. Ohmichi, S. Okubo, H. Ohta, K. Matsubayashi and Y. Uwatoko, *Appl. Magn. Reson.* **46** (2015) 987-992.
16. Zr-based bulk metallic glass as a cylinder material for high pressure apparatuses: K. Komatsu, K. Munakata, K. Matsubayashi, Y. Uwatoko, Y. Yokoyama, K. Sugiyama and M. Matsuda, *High Pressure Res.* **35** (2015) 254-262.
17. †Strange metal without magnetic criticality: T. Tomita, K. Kuga, Y. Uwatoko, P. Coleman and S. Nakatsuji, *Science* **349** (2015) 506-509.
18. Charge Transfer Induced Multifunctional Transitions with Sensitive Pressure Manipulation in a Metal–Organic Framework: J. Yang, L. Zhou, J. Cheng, Z. Hu, C. Kuo, C.-W. Pao, L. Jang, J.-F. Lee, J. Dai, S. Zhang, S. Feng, P. Kong, Z. Yuan, J. Yuan, Y. Uwatoko, T. Liu, C. Jin and Y. Long, *Inorg. Chem.* **54** (2015) 6433-6438.
19. †* Development of non-metallic diamond anvil cell and quantum oscillation measurement of CePt_2In_7 in a pulsed-magnet: A. Miyake, Y. Kohama, S. Ohta, Y. Hirose, R. Settai, K. Matsubayashi, Y. Uwatoko, A. Matsuo, K. Kindo and M. Tokunaga, *J. Phys.: Conf. Ser.* **592** (2015) 012149(1-6).
20. * High Pressure Measurements of the Resistivity of $\beta\text{-YbAlB}_4$: T. Tomita, K. Kuga, Y. Uwatoko and S. Nakatsuji, *J. Phys.: Conf. Ser.* **592** (2015) 012019.
21. †Development of multi-frequency ESR system for high-pressure measurements up to 2.5 GPa: T. Sakurai, K. Fujimoto, R. Matsui, K. Kawasaki, S. Okubo, H. Ohta, K. Matsubayashi, Y. Uwatoko and H. Tanaka, *J. Magn. Reson.* **259** (2015) 108-113.
22. †* Pressure-induced superconductivity in the iron-based ladder material BaFe_2S_3 : H. Takahashi, A. Sugimoto, Y. Nambu, T. Yamauchi, Y. Hirata, T. Kawakami, M. Avdeev, K. Matsubayashi, F. Du, C. Kawashima, H. Soeda, S. Nakano, Y. Uwatoko, Y. Ueda, T. J. Sato and K. Ohgushi, *Nature Mater.* **14** (2015) 1008-1012.
23. Charge disproportionation and the pressure-induced insulator–metal transition in cubic perovskite PbCrO_3 : J. Cheng, K. E. Kweon, S. A. Larregola, Y. Ding, Y. Shirako, L. G. Marshall, Z. -Y. Li, X. Li, A. M. D. Santos, M. R. Suchomel, K. Matsubayashi, Y. Uwatoko, G. S. Hwang, J. B. Goodenough and J. -S. Zhou, *Proc. Natl. Acad. Sci. U.S.A.* **112** (2015) 1670-1674.
24. Kondo Effect in CeX_c ($X_c = \text{S, Se, Te}$) Studied by Electrical Resistivity Measurements under High Pressure: Y. Hayashi, S. Takai, T. Matsumura, H. Tanida, M. Sera, K. Matsubayashi, Y. Uwatoko and A. Ochiai, *J. Phys. Soc. Jpn.* **85** (2016) 034704(1-7).
25. Superconducting and Fermi Surface Properties of Single Crystal Zr_2Co : A. Teruya, M. Kakihana, T. Takeuchi, D. Aoki, F. Honda, A. Nakamura, Y. Haga, K. Matsubayashi, Y. Uwatoko, H. Harima, M. Hedo, T. Nakama and Y. Onuki, *J. Phys. Soc. Jpn.* **85** (2016) 034706(1-10).
26. * Pressure dependence of the magnetic ground states in MnP: M. Matsuda, F. Ye, S. E. Dissanayake, J. -G. Cheng, S. Chi, J. Ma, H. D. Zhou, J. -Q. Yan, S. Kasamatsu, O. Sugino, T. Kato, K. Matsubayashi, T. Okada and Y. Uwatoko, *Phys. Rev. B* **93** (2016) 100405(1-5).
27. Quantum Criticality Beneath the Superconducting Dome in $\beta\text{-YbAlB}_4$: T. Tomita, K. Kuga, Y. Uwatoko and S. Nakatsuji, *J. Phys.: Conf. Ser.* **683** (2016) 012007(1-5).
28. Magnetic and Structural Properties of Metamagnetic $\text{MnCo}_{0.92}\text{Fe}_{0.08}\text{Ge}$ Compound: K. Ozono, Y. Mitsui, M. Hiroi, R. Y. Umetsu, K. Takahashi, K. Matsubayashi, Y. Uwatoko and K. Koyama, *Mater. Trans.* **57** (2016) 316-320.

* Joint research among groups within ISSP.

Osada group

We have systematically studied quantum Hall transport in monolayer-bilayer graphene heterojunctions on hexagonal boron nitride substrate. It has been found that the observed asymmetric transverse and Hall resistances across the junction are well understood by the Landauer-Büttiker edge transport picture assuming the pair annihilation of edge channels with opposite chirality at the junction. We show that the above picture works well even in the system where the monolayer and bilayer regions have different carrier density, which corresponds to the preceding work. This result demonstrates the bulk-edge correspondence at the boundary of two quantum Hall states on different crystal and band structures. In addition, we found that fine structures around charge neutrality points, which are considered to originate from the degeneracy breaking of zero-energy Landau levels of monolayer and bilayer graphene.

1. Edge State and Intrinsic Hole Doping in Bilayer Phosphorene: T. Osada, J. Phys. Soc. Jpn. **84** (2015) 013703(1-4).
2. Electronic Structure and the Properties of Phosphorene and Few-layer Black Phosphorus: S. Fukuoka, T. Taen and T. Osada, J. Phys. Soc. Jpn. **84** (2015) 121004(1-12).
3. Surface Transport in the $\nu = 0$ Quantum Hall Ferromagnetic State in the Organic Dirac Fermion System: T. Osada, J. Phys. Soc. Jpn. **84** (2015) 053704(1-4).

Yamashita group

As a joint project with Dr. Shishido at Osaka Prefecture University, we've performed dHvA measurements of the heavy-fermion superconductor CeCoIn₅. We succeeded to measure clear quantum oscillation measurements of the alpha bands of CeCoIn₅ down to 5 mK and up to 10 T. Unexpectedly, we found that the amplitude of the quantum oscillation is suppressed below 20 mK at 8 T (above H_{c2} of CeCoIn₅). At the same time, the frequency of the quantum oscillation shows a drop. These changes can be attributed to an emergence of a new ordered phase neighboring the superconducting phase. To clarify the property of the new phase, we started NMR measurements of CeCoIn₅ at ultra-low temperatures with Takigawa group, which is still a project under way in 2016. We've also measured longitudinal and transverse thermal transport measurements of a candidate material of quantum spin liquid, Ba₃CuSb₂O₉. We found that the longitudinal thermal conductivity is strongly suppressed in wide temperature range, showing that there are strong scatters of phonons. Further, a clear thermal Hall effect has been observed. Given that there are no mobile spin excitations in Ba₃CuSb₂O₉ from our measurements and NMR measurements, this thermal Hall effect is a phonon Hall effect. We believe that this work is the first systematic study of a thermal Hall effect of phonons including the temperature dependence of the thermal Hall conductivity. We've performed thermal Hall measurements of Mn₃Sn where the first observation of the anomalous Hall effect in antiferromagnetic system has been reported. A clear anomalous thermal Hall effect has been observed. We found that the temperature dependence of the Lorentz number implies that intrinsic scatterings play an important role for the anomalous Hall effect.

Materials Design and Characterization Laboratory

Hiroi group

One-Third Magnetization Plateau with a Preceding Novel Phase is discovered in Volborthite. We have synthesized high-quality single crystals of volborthite, and carried out high-field magnetization measurements up to 74 T and ⁵¹V NMR measurements up to 30 T. An extremely wide 1/3 magnetization plateau appears above 28 T and continues over 74 T at 1.4 K, which has not been observed in previous studies using polycrystalline samples. NMR spectra reveal an incommensurate order (most likely a spin-density wave order) below 22 T and a simple spin structure in the plateau phase. Moreover, a novel intermediate phase is found between 23 and 26 T, where the magnetization varies linearly with magnetic field and the NMR spectra indicate an inhomogeneous distribution of the internal magnetic field. This sequence of phases in volborthite bears a striking similarity to those of frustrated spin chains with a ferromagnetic nearest-neighbor coupling J_1 competing with an antiferromagnetic next-nearest-neighbor coupling J_2 . In addition, the metal-insulator transition (MIT) of VO₂ is discussed with particular emphasis on the structural instability of the rutile compounds toward dimerization. Ti substitution experiments reveal that the MIT is robust up to 20% Ti substitutions and occurs even in extremely thin V-rich lamellas in spinodally decomposed TiO₂-VO₂ composites, indicating that the MIT is insensitive to hole doping and essentially takes on a local character. These observations suggest that either electron correlation in the Mott-Hubbard sense or Peierls (Fermi-surface) instability plays a minor role on the MIT. Through a broad perspective of crystal chemistry on the rutile-related compounds, it is noted that VO₂ and another MIT compound NbO₂ in the family eventually lie just near the borderline between the two structural groups with the regular rutile structure and the distorted structures characterized by the formation of dimers with direct metal-metal bonding. It is also shown that the two compounds of the rutile form do not follow the general trends in structure observed for the other rutile compounds, giving clear evidence of an inherent structural instability present in the two compounds. The MITs of VO₂ and NbO₂ are natural consequences of structural transitions between the two groups, as all the d electrons are trapped in the bonding molecular

[†] Joint research with outside partners.

orbitals of dimers at low temperatures. Such dimer crystals are ubiquitously found in early transition metal compounds having chain-like structures, such as MoBr_3 , NbCl_4 , Ti_4O_7 , and V_4O_7 , the latter two of which also exhibit MITs probably of the same origin. In a broader sense, the dimer crystal is a kind of “molecular orbital crystals” in which virtual molecules made of transition metal atoms with partially-filled t_{2g} shells, such as dimers, trimers or larger ones, are generated by metal-metal bonding and are embedded into edge- or face-sharing octahedron networks of various kinds. The molecular orbital crystallization opens a natural route to stabilization of unpaired t_{2g} electrons in crystals.

1. ^{†*} Charge Order Induced by Cation Order in $\delta\text{-Ag}_{2/3}\text{V}_2\text{O}_5$: T. Baba, T. Yamauchi, S. Yamazaki, H. Ueda, M. Isobe, Y. Matsushita and Y. Ueda, *J. Phys. Soc. Jpn.* **84** (2015) 024718(1-7).
2. Large Diamagnetic Susceptibility from Petit Fermi Surfaces in $\text{LaV}_2\text{Al}_{20}$: T. Hirose, Y. Okamoto, J.-I. Yamaura and Z. Hiroi, *J. Phys. Soc. Jpn.* **84** (2015) 113701(1-5).
3. Magnetic Phase Diagram of the Breathing Pyrochlore Antiferromagnet $\text{LiGa}_{1-x}\text{In}_x\text{Cr}_4\text{O}_8$: Y. Okamoto, G. J. Nilsen, T. Nakazono and Z. Hiroi, *J. Phys. Soc. Jpn.* **84** (2015) 043707(1-5).
4. ^{*} Competing electronic states under pressure in the double-exchange ferromagnetic Peierls system $\text{K}_2\text{Cr}_8\text{O}_{16}$: T. Yamauchi, K. Hasegawa, H. Ueda, M. Isobe and Y. Ueda, *Phys. Rev. B* **92** (2015) 165115(1-7).
5. ^{*} Complex magnetostructural order in the frustrated spinel $\text{LiInCr}_4\text{O}_8$: G. J. Nilsen, Y. Okamoto, T. Masuda, J. Rodriguez-Carvajal, H. Mutka, T. Hansen and Z. Hiroi, *Phys. Rev. B* **91** (2015) 174435(1-8).
6. ^{†*} Effects of stoichiometry and substitution in quasi-one-dimensional iron chalcogenide BaFe_2S_3 : Y. Hirata, S. Maki, J.-I. Yamaura, T. Yamauchi and K. Ohgushi, *Phys. Rev. B* **92** (2015) 205109(1-7).
7. ^{†*} Structural stability of the Wadsley-type bronzes $\beta\text{-Ag}_{0.33}\text{V}_2\text{O}_5$ and $\beta\text{-Li}_{0.33}\text{V}_2\text{O}_5$ on compression: A breakdown of the two-leg ladder system in the nonsuperconducting high-pressure phase of $\beta\text{-Li}_{0.33}\text{V}_2\text{O}_5$: A. Grzechnik, Y. Ueda, T. Yamauchi, M. Hanfland, P. Hering, V. Potapkin and K. Frieze, *Phys. Rev. B* **91** (2015) 174113(1-8).
8. ^{*} Superconductivity in 122-type antimonide BaPtSb_2 : M. Imai, S. Ibuka, N. Kikugawa, T. Terashima, S. Uji, T. Yajima, H. Kageyama and I. Hase, *Phys. Rev. B* **91** (2015) 014513(1-8).
9. All-In–All-Out Magnetic Domains: X-Ray Diffraction Imaging and Magnetic Field Control: S. Tardif, S. Takeshita, H. Ohsumi, J.-I. Yamaura, D. Okuyama, Z. Hiroi, M. Takata and T.-H. Arima, *Phys. Rev. Lett.* **114** (2015) 147205.
10. ^{*} One-Third Magnetization Plateau with a Preceding Novel Phase in Volborthite: H. Ishikawa, M. Yoshida, K. Nawa, M. Jeong, S. Kramer, M. Horvatic, C. Berthier, M. Takigawa, M. Akaki, A. Miyake, M. Tokunaga, K. Kindo, J. Yamaura, Y. Okamoto and Z. Hiroi, *Phys. Rev. Lett.* **114** (2015) 227202(1-5).
11. ^{*} Pressure-induced Mott transition followed by a 24-K superconducting phase in BaFe_2S_3 : T. Yamauchi, Y. Hirata, Y. Ueda and K. Ohgushi, *Phys. Rev. Lett.* **115** (2015) 246402(1-5).
12. ^{†*} Semimetallic transport properties of epitaxially stabilized perovskite CaIrO_3 films: D. Hirai, J. Matsuno, D. Nishio-Hamane and H. Takagi, *Appl. Phys. Lett.* **107** (2015) 012104(1-4).
13. ^{*} Superconducting properties of $\text{BaTi}_2\text{Pn}_2\text{O}$ (Pn = Sb, Bi): T. Yajima, K. Nakano, Y. Nozaki and H. Kageyama, *Physica C* **504** (2015) 36-38.
14. $\text{Na}_2\text{MoO}_{2-\delta}\text{F}_{4+\delta}$ - a perovskite with a unique combination of atomic orderings and octahedral tilts: H. Ishikawa, I. Munaò, B. E. Bode, Z. Hiroi and P. Lightfoot, *Chem. Commun.* **51** (2015) 15469-15471.
15. ^{†*} Theoretical and experimental evidence for the post-cotunnite phase transition in zirconia at high pressure: D. Nishio-Hamane, H. Dekura, Y. Seto and T. Yagi, *Phys. Chem. Miner.* **42** (2015) 385-392.
16. ^{†*} Direct Observation of Short-Range Structural Coherence During a Charge Transfer Induced Spin Transition in a CoFe Prussian Blue Analogue by Transmission Electron Microscopy: M. Itoi, T. Jike, D. Nishio-Hamane, S. Udagawa, T. Tsuda, S. Kuwabata, K. Boukheddaden, M. J. Andrus and D. R. Talham, *J. Am. Chem. Soc.* **137** (2015) 14686-14693.
17. ^{†*} Effect of surfactant/water ratio and reagents' concentration on size distribution of manganese carbonate nanoparticles synthesized by microemulsion mediated route: G. Granata, F. Pagnanelli, D. Nishio-Hamane and T. Sasaki, *Appl. Surf. Sci.* **331** (2015) 463-471.
18. Superconductivity in LaPd_2As_2 with a collapsed 122 structure: S. Ganesanpotti, T. Yajima, K. Nakano, Y. Nozaki, T. Yamamoto, C. Tassel, Y. Kobayashi and H. Kageyama, *J. Alloys Compd.* **613** (2015) 370-374.
19. On the microscopic dynamics of the 'Einstein solids' $\text{AlV}_2\text{Al}_{20}$ and $\text{GaV}_2\text{Al}_{20}$, and of $\text{YV}_2\text{Al}_{20}$: a benchmark system for 'rattling' excitations: M. M. Koza, H. Mutka, Y. Okamoto, J.-I. Yamaura and Z. Hiroi, *Phys. Chem. Chem. Phys.* **17** (2015) 24837-24850.

* Joint research among groups within ISSP.

20. * Substrate-induced anion rearrangement in epitaxial thin films of $\text{LaSrCoO}_{4-x}\text{H}_x$: G. Bouilly, T. Yajima, T. Terashima, Y. Kususe, K. Fujita, C. Tassel, T. Yamamoto, K. Tanaka, Y. Kobayashi and H. Kageyama, *CrystEngComm* **16** (2015) 9669-9674.
21. †* Ferriakasaite-(La) and ferriandrosite-(La): new epidote-super group minerals from Ise, Mie Prefecture, Japan: M. Nagashima, D. Nishio-Hamane, N. Tomita, T. Minakawa and S. Inaba, *Mineral. Mag.* **79** (2015) 735.
22. * Imayoshiite, $\text{Ca}_3\text{Al}(\text{CO}_3)[\text{B}(\text{OH})_4](\text{OH})_6 \cdot 12\text{H}_2\text{O}$, a new mineral of ettringite group from Ise City, Mie Prefecture, Japan: D. Nishio-Hamane, M. Ohnishi, K. Momma, N. Shimobayashi, R. Miyawaki, T. Minakawa and S. Inaba, *Mineral. Mag.* **79** (2015) 413-423.
23. †* Pressure-induced superconductivity in the iron-based ladder material BaFe_2S_3 : H. Takahashi, A. Sugimoto, Y. Nambu, T. Yamauchi, Y. Hirata, T. Kawakami, M. Avdeev, K. Matsubayashi, F. Du, C. Kawashima, H. Soeda, S. Nakano, Y. Uwatoko, Y. Ueda, T. J. Sato and K. Ohgushi, *Nature Mater.* **14** (2015) 1008-1012.
24. Lifshitz metal-insulator transition induced by the all-in/all-out magnetic order in the pyrochlore oxide $\text{Cd}_2\text{Os}_2\text{O}_7$: Z. Hiroi, J. Yamaura, T. Hirose, I. Nagashima and Y. Okamoto, *APL Mater.* **3** (2015) 041501(1-11).
25. Spinodally decomposed nanostructures in a TiO_2 - VO_2 crystal: Z. Hiroi, T. Yoshida, J. Yamaura and Y. Okamoto, *APL Mater.* **3** (2015) 062508 (1-8).
26. * チタン酸塩化合物の高圧高温相転移: 浜根 大輔, *SPRING-8 利用研究成果集* **3** (2015) 6-9.
27. †* Impurity-Induced First-Order Phase Transitions in Highly Crystalline V_2O_3 Nanocrystals: Y. Ishiwata, E. Takahashi, K. Akashi, M. Imamura, J. Azuma, K. Takahashi, M. Kamada, H. Ishii, Y.-F. Liao, Y. Tezuka, Y. Inagaki, T. Kawae, D. Nishio-Hamane, M. Nantoh, K. Ishibashi and T. Kida, *Adv. Mater. Interfaces* **2** (2015) 1500132(1-6).
28. Structural instability of the rutile compounds and its relevance to the metal-insulator transition of VO_2 : Z. Hiroi, *Prog. Solid State Chem.* **43** (2015) 47-69.
29. †* Magnetic properties of $\text{Sm}_5\text{Fe}_{17}$ melt-spun ribbons and their borides: T. Saito and D. Nishio-Hamane, *AIMS Mat. Sci.* **2** (2015) 392-400.
30. †* Pressure-induced non-superconducting phase of $\beta\text{-Na}_{0.33}\text{V}_2\text{O}_5$ and the mechanism of high-pressure phase transitions in $\beta\text{-Na}_{0.33}\text{V}_2\text{O}_5$ and $\beta\text{-Li}_{0.33}\text{V}_2\text{O}_5$ at room temperature: A. Grzechnik, Y. Ueda, T. Yamauchi, M. Hanfland, P. Hering, V. Potapkin and K. Frieze, *J. Phys.: Condens. Matter* **28** (2016) 035401(1-9).
31. †* Hybrid Amine-Functionalized Graphene Oxide as a Robust Bifunctional Catalyst for Atmospheric Pressure Fixation of Carbon Dioxide using Cyclic Carbonates: V. B. Saptal, T. Sasaki, K. Harada, D. Nishio-Hamane and B. M. Bhanage, *ChemSusChem* **9** (2016) 644.
32. †* Light and SEM observation of opal phytoliths in the mulberry leaf: O. Tsutsui, R. Sakamoto, M. Obayashi, S. Yamakawa, T. Handa, D. Nishio-Hamane and I. Matsuda, *Flora* **218** (2016) 44-50.
33. 鉱物といふもの: 浜根 大輔, *物理学会誌* **70** (2015) 446-449.
34. * 新鉱物・宮久石の誕生について: 浜根 大輔, *岩石鉱物科学* **44** (2015) 57-59.

Kawashima group

We have been investigating quantum spin/boson systems and frustrated systems by means of large-scale numerical simulation. We also develop new numerical techniques. Our group's activities of 2015 include: (1) new interpretations and findings of computational results for the $\text{SU}(N)$ Heisenberg models with and without higher order interactions, (2) development of quantum Monte Carlo code for Bose systems in continuous media, (3) correlation between computational hardness of the spin glass instances and the thermodynamic properties and (3) large-scale non-equilibrium molecular dynamics simulation of bubble growth in under-pressured near-transition liquid.

1. Quantum Spin Liquid in Spin $1/2$ J_1 - J_2 Heisenberg Model on Square Lattice: Many-Variable Variational Monte Carlo Study Combined with Quantum-Number Projections: S. Morita, R. Kaneko and M. Imada, *J. Phys. Soc. Jpn.* **84** (2015) 024720(1-11).
2. Scaling relation for dangerously irrelevant symmetry-breaking fields: T. Okubo, K. Oshikawa, H. Watanabe and N. Kawashima, *Phys. Rev. B* **91** (2015) 174417(1-4).
3. $\text{SU}(N)$ Heisenberg model with multicolumn representations: T. Okubo, K. Harada, J. Lou and N. Kawashima, *Phys. Rev. B* **92** (2015) 1344048(1-5).

† Joint research with outside partners.

4. Thermal phase transition of generalized Heisenberg models for SU(N) spins on square and honeycomb lattices: T. Suzuki, K. Harada, H. Matsuo, S. Todo and N. Kawashima, *Phys. Rev. B* **91** (2015) 094414.
5. Variational Monte Carlo method in the presence of spin-orbit interaction and its application to Kitaev and Kitaev-Heisenberg models: M. Kurita, Y. Yamaji, S. Morita and M. Imada, *Phys. Rev. B* **92** (2015) 035122(1-11).
6. Thermal phase transitions to valence-bond-solid phase in the two dimensional; generalized SU(N) Heisenberg models: T. Suzuki, K. Harada, H. Matsuo, S. Todo and N. Kawashima, *J. Phys.: Conf. Ser.* **592** (2015) 012114.
7. Tensor network algorithm by coarse-graining tensor renormalization on finite periodic lattices: H.-H. Zhao, Z.-Y. Xie, T. Xiang and M. Imada, *Phys. Rev. B* **93** (2016) 125115(1-14).

Noguchi group

We have studied the membrane shape transformation in various conditions. We clarified the following behavior: (i) The absorption of banana-shaped proteins can induce polygonal membrane tubes and polyhedral vesicles. (2) Under chemical reaction an oil droplet can transform into vesicles via closing of a disk-like micelle. (3) At genus $g > 2$, the vesicle shape transformation from stomatocyte to discocyte is a discrete transition for low reduced volume.

1. [†]Formation of polyhedral vesicles and polygonal membrane tubes induced by banana-shaped proteins: H. Noguchi, *J. Chem. Phys.* **143** (2015) 243109.
2. Shape transitions of high-genus fluid vesicles: H. Noguchi, *Europhys. Lett.* **112** (2015) 58004.
3. Morphological changes of amphiphilic molecular assemblies induced by chemical reactions: K. M. Nakagawa and H. Noguchi, *Soft Matter* **11** (2015) 1403-1411.
4. Shape transformations of toroidal vesicles: H. Noguchi, A. Sakashita and M. Imai, *Soft Matter* **11** (2015) 193.
5. Shape deformation of lipid membranes by banana-shaped protein rods: Comparison with isotropic inclusions and membrane rupture: H. Noguchi, *Phys. Rev. E* **93** (2016) 052404(1-10).
6. [†]Monte Carlo study of the frame, fluctuation and internal tensions of fluctuating membranes with fixed area: H. Shiba, H. Noguchi and J.-B. Fournier, *Soft Matter* **12** (2016) 2373-2380.
7. Membrane tubule formation by banana-shaped proteins with or without transient network structure: H. Noguchi, *Sci. Rep.* **6** (2016) 20935.
8. Rheological evaluation of colloidal dispersions using the smoothed profile method: formulation and applications: J. J. Molina, K. Otomura, H. Shiba, H. Kobayashi, M. Sano and R. Yamamoto, *J. Fluid Mech.* **792** (2016) 590-619.

Materials Synthesis and Characterization group

1. ^{†*}Charge Order Induced by Cation Order in $\delta\text{-Ag}_{2/3}\text{V}_2\text{O}_5$: T. Baba, T. Yamauchi, S. Yamazaki, H. Ueda, M. Isobe, Y. Matsushita and Y. Ueda, *J. Phys. Soc. Jpn.* **84** (2015) 024718(1-7).
2. Magnetic properties of Sm-Fe-N bulk magnets prepared from $\text{Sm}_2\text{Fe}_{17}\text{N}_3$ melt-spun ribbons: T. Saito and D. Nishio-Hamane, *J. Appl. Phys.* **117** (2015) 17D130.
3. ^{†*}Effects of stoichiometry and substitution in quasi-one-dimensional iron chalcogenide BaFe_2S_3 : Y. Hirata, S. Maki, J.-I. Yamaura, T. Yamauchi and K. Ohgushi, *Phys. Rev. B* **92** (2015) 205109(1-7).
4. ^{†*}Structural stability of the Wadsley-type bronzes $\beta\text{-Ag}_{0.33}\text{V}_2\text{O}_5$ and $\beta\text{-Li}_{0.33}\text{V}_2\text{O}_5$ on compression: A breakdown of the two-leg ladder system in the nonsuperconducting high-pressure phase of $\beta\text{-Li}_{0.33}\text{V}_2\text{O}_5$: A. Grzechnik, Y. Ueda, T. Yamauchi, M. Hanfland, P. Hering, V. Potapkin and K. Frieze, *Phys. Rev. B* **91** (2015) 174113(1-8).
5. ^{*}Superconductivity in 122-type antimonide BaPt_2Sb_2 : M. Imai, S. Ibuka, N. Kikugawa, T. Terashima, S. Uji, T. Yajima, H. Kageyama and I. Hase, *Phys. Rev. B* **91** (2015) 014513(1-8).
6. ^{†*}Semimetallic transport properties of epitaxially stabilized perovskite CaIrO_3 films: D. Hirai, J. Matsuno, D. Nishio-Hamane and H. Takagi, *Appl. Phys. Lett.* **107** (2015) 012104(1-4).
7. ^{*}Superconducting properties of $\text{BaTi}_2\text{Pn}_2\text{O}$ (Pn = Sb, Bi): T. Yajima, K. Nakano, Y. Nozaki and H. Kageyama, *Physica C* **504** (2015) 36-38.

* Joint research among groups within ISSP.

8. ^{†*}Theoretical and experimental evidence for the post-cotunnite phase transition in zirconia at high pressure: D. Nishio-Hamane, H. Dekura, Y. Seto and T. Yagi, *Phys. Chem. Miner.* **42** (2015) 385-392.
9. ^{†*}Direct Observation of Short-Range Structural Coherence During a Charge Transfer Induced Spin Transition in a CoFe Prussian Blue Analogue by Transmission Electron Microscopy: M. Itoi, T. Jike, D. Nishio-Hamane, S. Udagawa, T. Tsuda, S. Kuwabata, K. Boukheddaden, M. J. Andrus and D. R. Talham, *J. Am. Chem. Soc.* **137** (2015) 14686-14693.
10. ^{†*}Effect of surfactant/water ratio and reagents' concentration on size distribution of manganese carbonate nanoparticles synthesized by microemulsion mediated route: G. Granata, F. Pagnanelli, D. Nishio-Hamane and T. Sasaki, *Appl. Surf. Sci.* **331** (2015) 463-471.
11. New hard magnetic phase in Mn–Ga–Al system alloys: T. Saito and D. Nishio-Hamane, *J. Alloys Compd.* **632** (2015) 486.
12. ^{*}Substrate-induced anion rearrangement in epitaxial thin films of $\text{LaSrCoO}_{4-x}\text{H}_x$: G. Bouilly, T. Yajima, T. Terashima, Y. Kususe, K. Fujita, C. Tassel, T. Yamamoto, K. Tanaka, Y. Kobayashi and H. Kageyama, *CrystEngComm* **16** (2015) 9669-9674.
13. ^{†*}Ferriakasaite-(La) and ferriandrosite-(La): new epidote-supergroup minerals from Ise, Mie Prefecture, Japan: M. Nagashima, D. Nishio-Hamane, N. Tomita, T. Minakawa and S. Inaba, *Mineral. Mag.* **79** (2015) 735.
14. ^{*}Imayoshiite, $\text{Ca}_3\text{Al}(\text{CO}_3)[\text{B}(\text{OH})_4](\text{OH})_6 \cdot 12\text{H}_2\text{O}$, a new mineral of ettringite group from Ise City, Mie Prefecture, Japan: D. Nishio-Hamane, M. Ohnishi, K. Momma, N. Shimobayashi, R. Miyawaki, T. Minakawa and S. Inaba, *Mineral. Mag.* **79** (2015) 413-423.
15. ^{†*}Pressure-induced superconductivity in the iron-based ladder material BaFe_2S_3 : H. Takahashi, A. Sugimoto, Y. Nambu, T. Yamauchi, Y. Hirata, T. Kawakami, M. Avdeev, K. Matsubayashi, F. Du, C. Kawashima, H. Soeda, S. Nakano, Y. Uwatoko, Y. Ueda, T. J. Sato and K. Ohgushi, *Nature Mater.* **14** (2015) 1008-1012.
16. Synthesis of oxamate and urea by oxidative single and double carbonylation of amines using immobilized palladium metal-containing ionic liquid@SBA-15: S. T. Gadge, E. N. Kusumawati, K. Harada, T. Sasaki, D. Nishio-Hamane and B. M. Bhanage, *J. Mol. Catal. A: Chem.* **400** (2015) 170.
17. Synthesis of Polyester Amide by Carbonylation–Polycondensation Reaction Using Immobilized Palladium Metal Containing Ionic Liquid on SBA-15 as a Phosphine-Free Catalytic System: A. Satapathy, S. T. Gadge, E. N. Kusumawati, K. Harada, T. Sasaki, D. Nishio-Hamane and B. M. Bhanage, *Catal. Lett.* **145** (2015) 824.
18. ^{*}チタン酸塩化合物の高圧高温相転移：浜根 大輔，SPRING-8 利用研究成果集 **3** (2015) 6-9.
19. ^{†*}Impurity-Induced First-Order Phase Transitions in Highly Crystalline V_2O_3 Nanocrystals: Y. Ishiwata, E. Takahashi, K. Akashi, M. Imamura, J. Azuma, K. Takahashi, M. Kamada, H. Ishii, Y.-F. Liao, Y. Tezuka, Y. Inagaki, T. Kawae, D. Nishio-Hamane, M. Nantoh, K. Ishibashi and T. Kida, *Adv. Mater. Interfaces* **2** (2015) 1500132(1-6).
20. ^{†*}Magnetic properties of $\text{Sm}_5\text{Fe}_{17}$ melt-spun ribbons and their borides: T. Saito and D. Nishio-Hamane, *AIMS Mat. Sci.* **2** (2015) 392-400.
21. ^{†*}Pressure-induced non-superconducting phase of $\beta\text{-Na}_{0.33}\text{V}_2\text{O}_5$ and the mechanism of high-pressure phase transitions in $\beta\text{-Na}_{0.33}\text{V}_2\text{O}_5$ and $\beta\text{-Li}_{0.33}\text{V}_2\text{O}_5$ at room temperature: A. Grzechnik, Y. Ueda, T. Yamauchi, M. Hanfland, P. Hering, V. Potapkin and K. Frieze, *J. Phys.: Condens. Matter* **28** (2016) 035401(1-9).
22. ^{†*}Hybrid Amine-Functionalized Graphene Oxide as a Robust Bifunctional Catalyst for Atmospheric Pressure Fixation of Carbon Dioxide using Cyclic Carbonates: V. B. Saptal, T. Sasaki, K. Harada, D. Nishio-Hamane and B. M. Bhanage, *ChemSusChem* **9** (2016) 644.
23. ^{†*}Light and SEM observation of opal phytoliths in the mulberry leaf: O. Tsutsui, R. Sakamoto, M. Obayashi, S. Yamakawa, T. Handa, D. Nishio-Hamane and I. Matsuda, *Flora* **218** (2016) 44-50.
24. ^{*}新鉱物・宮久石の誕生について：浜根 大輔，岩石鉱物科学 **44** (2015) 57-59.

[†] Joint research with outside partners.

Shibayama group

Shibayama group has been exploring the structure and dynamics of soft matter, especially polymer gels, micelles, and phenolic resin, utilizing a combination of small-angle neutron scattering (SANS), small-angle X-ray scattering (SAXS), and dynamic light scattering (DLS). The objectives are to elucidate the relationship between the structure and variety of novel properties/functions of polymer gels/resins. The highlights of 2015 include (1) development of high-toughness Ion gel for CO₂ separation, (2) gelation and cross-link inhomogeneity of phenolic resins, (3) structure evolution of catalyst ink for fuel cell in drying process, (4) rubber elasticity for percolation network consisting of Gaussian chains, (5) gelation mechanism of Tetra-Armed Poly(ethylene glycol) in aprotic ionic liquid, (6) structural analysis of lipophilic polyelectrolyte solutions and gels in low-polar solvents and so on.

1. Gelation Mechanism of Tetra-Armed Poly(ethylene glycol) in Aprotic Ionic Liquid Containing Non-volatile Proton Source, Protic Ionic Liquid: K. Hashimoto, K. Fujii, K. Nishi, T. Sakai, N. Yoshimoto, M. Morita and M. Shibayama, *J. Phys. Chem. B* **119** (2015) 4795-4801.
2. Microscopic Solvation Structure of Glucose in 1-Ethyl-3-methylimidazolium Methylphosphonate Ionic Liquid: K. Hirose, K. Fujii, K. Hashimoto, Y. Umebayashi and M. Shibayama, *J. Phys. Chem. B* **119** (2015) 6262.
3. Electrophoretic mobility of semi-flexible double-stranded DNA in defect-controlled polymer networks: Mechanism investigation and role of structural parameters: K. Khairulina, X. Li, K. Nishi, M. Shibayama, U.-I. Chung and T. Sakai, *J. Chem. Phys.* **142** (2015) 234904.
4. Rubber elasticity for percolation network consisting of Gaussian chains: K. Nishi, H. Noguchi, T. Sakai and M. Shibayama, *J. Chem. Phys.* **143** (2015) 184905.
5. Phase Behavior of Block Copolymers in Selective Supercritical Solvent: M. M. M. Ito, K. Ito, M. Shibayama, K. Sugiyama and H. Yokoyama, *Macromolecules* **48** (2015) 3590.
6. Structural Analysis of Lipophilic Polyelectrolyte Solutions and Gels in Low-Polar Solvents: K. Nishi, S. Tochioka, T. Hiroi, T. Yamada, K. Kokado, T.-H. Kim, E. P. Gilbert, K. Sada and M. Shibayama, *Macromolecules* **48** (2015) 3613.
7. 高分子ゲルおよびミセルの中性子散乱に魅せられて：柴山 充弘，波紋 **25** (2015) 120-125.
8. Carbon Dioxide Separation Using a High-toughness Ion Gel with a Tetra-armed Polymer Network: K. Fujii, T. Makino, K. Hashimoto, T. Sakai, M. Kanakubo and M. Shibayama, *Chem. Lett.* **44** (2015) 17.
9. Reliable Hydrogel with Mechanical “Fuse Link” in an Aqueous Environment: S. Kondo, T. Hiroi, Y.-S. Han, T.-H. Kim, M. Shibayama, U.-I. Chung and T. Sakai, *Adv. Mater.* **27** (2015) 7407.
10. Dynamic light scattering study on curing mechanisms of novolac-type phenolic resins: Y. Shudo, A. Izumi, T. Takeuchi, T. Nakao and M. Shibayama, *Polym. J.* **47** (2015) 79-81.
11. Gelation Kinetics and Polymer Network Dynamics of Homogeneous Tetra-PEG Gels: T. Birshstein, *Macromol. Symp.* **348** (2015) 9.
12. Supercoiling transformation of chemical gels: M. Asai, T. Katashima, T. Sakai and M. Shibayama, *Soft Matter* **11** (2015) 7101.
13. Multiblock copolymers exhibiting spatio-temporal structure with autonomous viscosity oscillation: M. Onoda, T. Ueki, M. Shibayama and R. Yoshida, *Sci. Rep.* **5** (2015) 15792.
14. Structural evolution of a catalyst ink for fuel cells during the drying process investigated by CV-SANS: T. Kusano, T. Hiroi, K. Amemiya, M. Ando, T. Takahashi and M. Shibayama, *Polym. J.* **47** (2015) 546.
15. Self-oscillating AB diblock copolymer developed by post modification strategy: T. Ueki, M. Onoda, R. Tamate, M. Shibayama and R. Yoshida, *Chaos* **25** (2015) 064605.
16. Nearly Ideal Polymer Network Ion Gel Prepared in pH-Buffering Ionic Liquid: K. Hashimoto, K. Fujii, K. Nishi, T. Sakai and M. Shibayama, *Macromolecules* **49** (2016) 344.
17. Transitions of Aggregation States for Concentrated Carbon Nanotube Dispersion: T. Hiroi, S. Ata and M. Shibayama, *J. Phys. Chem. C* **120** (2016) 5776.
18. 小角 X 線散乱法によるフェノール樹脂ベークン過程のその場解析：和泉 篤士，若林みどり，首藤 靖幸，中尾 俊夫，柴山 充弘，ネットワークポリマー **2015** (2015) 83-88.

* Joint research among groups within ISSP.

19. Small-angle Neutron Scattering of Polysaccharide Hydrogels: M. Shibayama, in: *Polysaccharide hydrogels: Characterization and Biomedical Applications*, Ch 7, edited by P. Matricardi, F. Alhaique and T. Coviello, (Pan Stanford Publishing Pte. Ltd., Singapore, 2016), 245-264.

Yoshizawa group

A systematic study on spin dynamics in a two-dimensional transition-metal Ni oxide has been carried out with use of the high resolution chopper spectrometer installed at BL12 in the Material and Life Science Facility, J-PARC. The checkerboard-type spin-charge ordering in the highly hole-doped region of the layered nickelate was studied in detail. The nature of the excitation spectra and the thermodynamic properties in the checkerboard phase was found to show qualitative differences from those in the stripe phase. Magnetic properties of a family of Ce-based non-centrosymmetric heavy fermion compounds CeTSi_3 (T=transition metal ions) were also studied.

1. Neutron Diffraction Study of Parasitic Nd-Moment Order in the Checkerboard-Type Phase $\text{Nd}_{1.3}\text{Sr}_{0.7}\text{NiO}_4$: R. Kobayashi, H. Yoshizawa, M. Matsuda, R. Kajimoto, K. Ishizaka and Y. Tokura, *J. Phys. Soc. Jpn.* **84** (2015) 064711.
2. Resistance Anomalies Accompanying Crossover from Heavy-Fermion Regime to Intermediate-Valence Regime: A Study of Cu–Ni Substitution and Pressure Effects on CeCu_2Si_2 : Y. Ikeda, Y. Ito, S. Araki, T. C. Kobayashi and H. Yoshizawa, *J. Phys. Soc. Jpn.* **84** (2015) 024702.
3. Single-Crystal Neutron Diffraction Study of the Heavy-Electron Superconductor CeNiGe_3 : Y. Ikeda, D. Ueta, H. Yoshizawa, A. Nakao, K. Munakata and T. Ohhara, *J. Phys. Soc. Jpn.* **84** (2015) 123701.
4. Transport and Thermodynamic Studies of Stripe and Checkerboard Ordering in Layered Nickel Oxides $\text{R}_{2-x}\text{Sr}_x\text{NiO}_4$ (R = La and Nd): Y. Ikeda, S. Suzuki, T. Nakabayashi, H. Yoshizawa, T. Yokoo and S. Itoh, *J. Phys. Soc. Jpn.* **84** (2015) 023706.
5. [†]*Pauli-limited superconductivity and antiferromagnetism in the heavy-fermion compound $\text{CeCo}(\text{In}_{1-x}\text{Zn}_x)_5$: M. Yokoyama, H. Mashiko, R. Otaka, Y. Sakon, K. Fujimura, K. Tenya, A. Kondo, K. Kindo, Y. Ikeda, H. Yoshizawa, Y. Shimizu, Y. Kono and T. Sakakibara, *Phys. Rev. B* **92** (2015) 184509(9).
6. *Temperature and composition phase diagram in the iron-based ladder compounds $\text{Ba}_{1-x}\text{Cs}_x\text{Fe}_2\text{Se}_3$: T. Hawaii, Y. Nambu, K. Ohgushi, F. Du, Y. Hirata, M. Avdeev, Y. Uwatoko, Y. Sekine, H. Fukazawa, J. Ma, S. Chi, Y. Ueda, H. Yoshizawa and T. J. Sato, *Phys. Rev. B* **91** (2015) 184416.
7. Characterization of Ferromagnetic Order in CePd_2P_2 : Y. Ikeda, H. Yoshizawa, S. Konishi, S. Araki, T. C. Kobayashi, T. Yokoo and S. Itoh, *J. Phys.: Conf. Ser.* **592** (2015) 012013.
8. *Science from the Initial Operation of HRC: S. Itoh, T. Yokoo, T. Masuda, H. Yoshizawa, M. Soda, Y. Ikeda, S. Ibuka, D. Kawana, T. J. Sato, Y. Nambu, K. Kuwahara, S.-I. Yano, J. Akimitsu, Y. Kaneko, Y. Tokura, M. Fujita, M. Hase, K. Iwasa, H. Hiraka, T. Fukuda, K. Ikeuchi, K. Yoshida, T. Yamaguchi, K. Ono and Y. Endoh, *JPS Conf. Proc.* **8** (2015) 034001.
9. Inelastic Neutron Scattering Study of Stripe and Overdoped Checkerboard Ordering in Layered Nickel Oxide $\text{Nd}_{2-x}\text{Sr}_x\text{NiO}_4$: Y. Ikeda, S. Suzuki, T. Nakabayashi, H. Yoshizawa, T. Yokoo and S. Itoh, *J. Phys. Soc. Jpn.* **85** (2016) 023701.

Yamamuro group

Our laboratory is studying chemical physics of complex condensed matters by using neutron scattering, X-ray diffraction, calorimetric, dielectric, and viscoelastic techniques. Our target materials are glasses, liquids, and various disordered systems. This year, we have succeeded to observe the inelastic neutron scattering spectra of palladium hydride nanoparticles. The peaks characteristic to the nanoparticles are attributed to the vibration of the H atoms located at the tetrahedral sites of the Pd lattice, which were predicted by our neutron powder diffraction work. Another topic is on the dynamics of a reverse osmotic membrane consisting of aromatic polyamide and water. This system is remarked for seawater desalination and waste-water reclamation. Several quasielastic neutron scattering experiments in a wide time range (0.5 ps to 5 ns) revealed that the polyamide network is drastically plasticized by water and moving with water even at 240 K. This information will give a new insight into the mechanism of water purification. Other than above topics, we have made some progresses in the studies on vapor-deposited molecular glasses and ionic liquids with plastically crystalline phases.

1. Glass transition and positional ordering of hydrogen in bulk and nanocrystalline palladium: H. Akiba, H. Kobayashi, H. Kitagawa, M. Kofu and O. Yamamuro, *Phys. Rev. B* **92** (2015) 064202.

[†] Joint research with outside partners.

2. Thermal and Structural Studies of Imidazolium-Based Ionic Liquids with and without Liquid-Crystalline Phases: The Origin of Nanostructure: F. Nemoto, M. Kofu and O. Yamamuro, *J. Phys. Chem. B* **119** (2015) 5028-5034.
3. Quasielastic neutron scattering studies on glass-forming ionic liquids with imidazolium cations: M. Kofu, M. Tyagi, Y. Inamura, K. Miyazaki and O. Yamamuro, *J. Chem. Phys.* **143** (2015) 234502.
4. Inelastic neutron scattering study on boson peaks of imidazolium-based ionic liquids: M. Kofu, Y. Inamura, Y. Moriya, A. Podlesnyak, G. Ehlers and O. Yamamuro, *Journal of Molecular Liquids* **210B** (2015) 164-168.
5. Connecting thermodynamics and dynamics in a supercooled liquid: Cresolphthalein-dimethylether: S. Samanta, O. Yamamuro and R. Richert, *Thermochimica Acta* **636** (2016) 57-62.
6. Effect of water on the structure of a prototype ionic liquid: O. Borodin, D. L. Price, B. Aoun, M. A. González, J. B. Hooper, M. Kofu, S. Kohara, O. Yamamuro and M.-L. Saboungi, *Phys. Chem. Chem. Phys.* (2016) 10.1039/C6CP02191C, in print.
7. 中性子散乱で観たイオン液体の階層的構造とダイナミクス: 山室 修, 根本 文也, 古府 麻衣子, 高圧力の科学と技術 **25** (2015) 200-207.
8. 低温蒸着法で作製した単純分子ガラスの構造: 山室 修, 水野 勇希, 古府 麻衣子, 日本結晶学会誌 **58** (2016) 13-17.
9. イミダゾリウム系イオン液体の階層的・ガラスダイナミクス: 古府 麻衣子, 山室 修, 日本結晶学会誌 **58** (2016) 18-23.

Masuda group

The goal of our research is to discover a new quantum phenomenon and to reveal the mechanism of it. In this fiscal year we studied the following topics; complex magnetostructural order in the frustrated spinel $\text{LiInCr}_4\text{O}_8$, magnetic model in multiferroic $\text{NdFe}_3(\text{BO}_3)_4$, specific heats of triangular spin tube in magnetic fields, and magnetic anti-cancer compound for magnet-guided delivery and magnetic resonance imaging.

1. *Complex magnetostructural order in the frustrated spinel $\text{LiInCr}_4\text{O}_8$: G. J. Nilsen, Y. Okamoto, T. Masuda, J. Rodriguez-Carvajal, H. Mutka, T. Hansen and Z. Hiroi, *Phys. Rev. B* **91** (2015) 174435(1-8).
2. Magnetic model in multiferroic $\text{NdFe}_3(\text{BO}_3)_4$ investigated by inelastic neutron scattering: S. Hayashida, M. Soda, S. Itoh, T. Yokoo, K. Ohgushi, D. Kawana, H. M. Rønnow and T. Masuda, *Phys. Rev. B* **92** (2015) 054402.
3. マルチフェロイック物質 $\text{Ba}_2\text{CoGe}_2\text{O}_7$ におけるスピン・ネマティック相互作用の観測: 益田 隆嗣, 左右田 稔, 固体物理 **50** (2015) 111.
4. Inelastic Neutron Scattering on Multiferroics $\text{NdFe}_3(\text{BO}_3)_4$: S. Hayashida, M. Soda, S. Itoh, T. Yokoo, K. Ohgushi, D. Kawana and T. Masuda, *Physics Procedia* **75** (2015) 127.
5. Specific Heats of Triangular Spin Tube in Magnetic Fields: H. Manaka, M. Hagihara, S. Hayashida, M. Soda, T. Masuda and Y. Miura, *Physics Procedia* **75** (2015) 718.
6. A magnetic anti-cancer compound for magnet-guided delivery and magnetic resonance imaging: H. Eguchi, M. Umemura, R. Kurotani, H. Fukumura, I. Sato, J.-H. Kim, Y. Hoshino, J. Lee, N. Amemiya, M. Sato, K. Hirata, D. J. Singh, T. Masuda, M. Yamamoto, T. Urano, K. Yoshida, K. Tanigaki, M. Yamamoto, M. Sato, S. Inoue, I. Aoki and Y. Ishikawa, *Sci. Rep.* **5** (2015) 9194.
7. *Science from the Initial Operation of HRC: S. Itoh, T. Yokoo, T. Masuda, H. Yoshizawa, M. Soda, Y. Ikeda, S. Ibuka, D. Kawana, T. J. Sato, Y. Nambu, K. Kuwahara, S.-I. Yano, J. Akimitsu, Y. Kaneko, Y. Tokura, M. Fujita, M. Hase, K. Iwasa, H. Hiraka, T. Fukuda, K. Ikeuchi, K. Yoshida, T. Yamaguchi, K. Ono and Y. Endoh, *JPS Conf. Proc.* **8** (2015) 034001.
8. Crystal Field Excitations in the Breathing Pyrochlore Antiferromagnet $\text{Ba}_3\text{Yb}_2\text{Zn}_5\text{O}_{11}$: T. Haku, M. Soda, M. Sera, K. Kimura, S. Itoh, T. Yokoo and T. Masuda, *J. Phys. Soc. Jpn.* **85** (2016) 034721.
9. Dielectric and Magnetic Properties in Relaxor Magnet LuFeCoO_4 : M. Soda and T. Masuda, *J. Phys. Soc. Jpn.* **85** (2016) 034713.
10. Spin Model of O_2 -Based Magnet in a Nanoporous Metal Complex: M. Soda, Y. Honma, S. Takamizawa, S. Ohira-Kawamura, K. Nakajima and T. Masuda, *J. Phys. Soc. Jpn.* **85** (2016) 034717.
11. Magnetic ordering of the buckled honeycomb lattice antiferromagnet $\text{Ba}_2\text{NiTeO}_6$: S. Asai, M. Soda, K. Kasatani, T. Ono, M. Avdeev and T. Masuda, *Phys. Rev. B* **93** (2016) 024412.

* Joint research among groups within ISSP.

International MegaGauss Science Laboratory

Takeyama group

Single-walled carbon nanotubes (SWNT) with a typical tube diameter an order of 1 nm are characterized by the quasi-one dimensional system. Optical properties are enriched by the enhancement of the excitonic band-edge structure. When a magnetic flux penetrates through the tube, the electronic band structure is subjected to substantial modulation via the mechanism of Aharonov-Bohm (A-B) effect. This effect lifts the quantum degeneracy of the band-edge states, induces splittings of the exciton optical absorption peak in megagauss magnetic field. The magneto-optical A-B splitting was investigated in magnetic fields of up to 300 – 400 T by observing band-edge exciton absorption spectra. The spectra evolved as a simple bright and dark exciton splitting with a linear magnetic field dependence as has been predicted by T. Ando. The ratio of the splitting is governed by a microscopic information of the environment dielectric substances surrounding the SWNTs.

1. *Magnetic-Field-Induced Insulator–Metal Transition in $(\text{Pr}_{1-y}\text{Y}_y)_{0.7}\text{Ca}_{0.3}\text{CoO}_3$ at Ultrahigh Magnetic Fields: S. Lee, Y. H. Matsuda, T. Naito, D. Nakamura and S. Takeyama, *J. Phys. Soc. Jpn.* **84** (2015) 044703-7.
2. Exciton splitting in semiconducting carbon nanotubes in ultrahigh magnetic fields above 300 T: D. Nakamura, T. Sasaki, W. Zhou, H. Liu, H. Kataura and S. Takeyama, *Phys. Rev. B* **91** (2015) 235427.
3. †*Phase boundary of θ phase of solid oxygen in ultrahigh magnetic fields: T. Nomura, Y. H. Matsuda, S. Takeyama, A. Matsuo, K. Kindo and T. C. Kobayashi, *Phys. Rev. B* **92** (2015) 064109 (4 pages).
4. †*One-Way Transparency of Light in Multiferroic CuB_2O_4 : S. Toyoda, N. Abe, S. Kimura, Y. H. Matsuda, T. Nomura, A. Ikeda, S. Takeyama and T. Arima, *Phys. Rev. Lett.* **115** (2015) 267207 (5 pages).
5. *Effect of very high magnetic field on the optical properties of firefly light emitter oxyluciferin: W. Zhou, D. Nakamura, Y. Wang, T. Mochizuki, H. Akiyama and S. Takeyama, *J. Lumin.* **165** (2015) 15.
6. †*Anomalous diamagnetic shifts in InP-GaP lateral quantum-wires: Y. H. Shin, B. K. Choi, Y. Kim, J. D. Song, D. Nakamura, Y. H. Matsuda and S. Takeyama, *Opt. Express* **23** (2015) 28349-28357.

Kindo group

Long pulsed magnetic field has been improved. The maximum field of long pulse magnet was limited to 36 T due to the control system. We have improved the control system and succeeded in generating 42.5 T. The other improvement is development of almost perfect flat-top field. We have added a small coil into the bore of the large long pulse magnet and controlled the small coil to obtain the flat-top magnetic field. We have succeeded in generating 40 ± 0.005 T for 0.2 sec. These improvements will expand the possibility of the high field study on the condensed matter physics.

1. *Field Evolution of Quantum Critical and Heavy Fermi-Liquid Components in the Magnetization of the Mixed Valence Compound β -YbAlB₄: Y. Matsumoto, K. Kuga, Y. Karaki, Y. Shimura, T. Sakakibara, M. Tokunaga, K. Kindo and S. Nakatsuji, *J. Phys. Soc. Jpn.* **84** (2015) 024710(1-7).
2. *Possible Excitonic Phase of Graphite in the Quantum Limit State: K. Akiba, A. Miyake, H. Yaguchi, A. Matsuo, K. Kindo and M. Tokunaga, *J. Phys. Soc. Jpn.* **84** (2015) 054709(1-6).
3. †Spin-Liquid Ground State in the Spin 1/2 Distorted Diamond Chain Compound $\text{K}_3\text{Cu}_3\text{AlO}_2(\text{SO}_4)_4$: M. Fujihala, H. Koorikawa, S. Mitsuda, M. Hagihala, H. Morodomi, T. Kawae, A. Matsuo and K. Kindo, *J. Phys. Soc. Jpn.* **84** (2015) 073702(1-4).
4. *X-ray Absorption Spectroscopy in the Heavy Fermion Compound α -YbAlB₄ at High Magnetic Fields: T. T. Terashima, Y. H. Matsuda, K. Kuga, S. Suzuki, Y. Matsumoto, S. Nakatsuji, A. Kondo, K. Kindo, N. Kawamura, M. Mizumaki and T. Inami, *J. Phys. Soc. Jpn.* **84** (2015) 114715(1-4).
5. †High-magnetic-field phase transitions and H–T phase diagram of the Kagome-staircase compound $\text{Ni}_3\text{V}_2\text{O}_8$: Z. Q. Lin, M. Yang, H. W. Wang, Q. Guo, Y. J. Liu, X. T. Han, Y. B. Han, J. F. Wang, Z. Z. He and K. Kindo, *J. Magn. Magn. Mater.* **382** (2015) 7.
6. †Evolution of exchange interaction constants across magnetic phase transitions in the chromium spinel oxide CdCr_2O_4 : S. Kimura, Y. Sawada, Y. Narumi, K. Watanabe, M. Hagiwara, K. Kindo and H. Ueda, *Phys. Rev. B* **92** (2015) 144410(1-9).
7. †*Pauli-limited superconductivity and antiferromagnetism in the heavy-fermion compound $\text{CeCo}(\text{In}_{1-x}\text{Zn}_x)_5$: M. Yokoyama, H. Mashiko, R. Otaka, Y. Sakon, K. Fujimura, K. Tenya, A. Kondo, K. Kindo, Y. Ikeda, H. Yoshizawa, Y. Shimizu, Y. Kono and T. Sakakibara, *Phys. Rev. B* **92** (2015) 184509(1-9).

† Joint research with outside partners.

8. ^{†*}Phase boundary of θ phase of solid oxygen in ultrahigh magnetic fields: T. Nomura, Y. H. Matsuda, S. Takeyama, A. Matsuo, K. Kindo and T. C. Kobayashi, *Phys. Rev. B* **92** (2015) 064109 (4 pages).
9. [†]Successive magnetic phase transitions in α -RuCl₃: XY-like frustrated magnet on the honeycomb lattice: Y. Kubota, H. Tanaka, T. Ono, Y. Narumi and K. Kindo, *Phys. Rev. B* **91** (2015) 094422.
10. [†]Valence-specific magnetization of the charge-ordered multiferroelectric LuFe₂O₄ using soft x-ray magnetic circular dichroism under 30 T pulsed high magnetic fields: Y. Narumi, T. Nakamura, K. Saito, T. Morioka, Y. Fukada, T. Kambe, N. Ikeda, Y. Kotani, T. Kinoshita, K. Kindo and H. Nojiri, *Phys. Rev. B* **91** (2015) 014410.
11. Generation of flat-top pulsed magnetic fields with feedback control approach: Y. Kohama and K. Kindo, *Rev. Sci. Instrum.* **86** (2015) 104701(5).
12. ^{*}One-Third Magnetization Plateau with a Preceding Novel Phase in Volborthite: H. Ishikawa, M. Yoshida, K. Nawa, M. Jeong, S. Kramer, M. Horvatic, C. Berthier, M. Takigawa, M. Akaki, A. Miyake, M. Tokunaga, K. Kindo, J. Yamaura, Y. Okamoto and Z. Hiroi, *Phys. Rev. Lett.* **114** (2015) 227202(1-5).
13. ^{*}パルス強磁場下における磁気熱量効果／比熱測定手法の開発：木原 工，小濱 芳允，徳永 将史，金道 浩一，*固体物理* **50** (2015) 371.
14. ^{†*}Anisotropy in the upper critical field of FeSe and FeSe_{0.33}Te_{0.67} single crystals: J. L. Her, Y. Kohama, Y. H. Matsuda, K. Kindo, W.-H. Yang, D. A. Chareev, E. S. Mitrofanova, O. S. Volkova, A. N. Vasiliev and J.-Y. Lin, *Supercond. Sci. Technol.* **28** (2015) 045013 (6 pages).
15. [†]Large magnetoresistance of EuPtP_{1-x}As_x: M. Sugishima, H. Wada, A. Mitsuda, A. Kondo and K. Kindo, *Phys. Status Solidi B* **252** (2015) 2784-2788.
16. ^{†*}Superconductivity protected by spin-valley locking in ion-gated MoS₂: Y. Saito, Y. Nakamura, M. S. Bahramy, Y. Kohama, J. Ye, Y. Kasahara, Y. Nakagawa, M. Onga, M. Tokunaga, T. Nojima, Y. Yanase and Y. Iwasa, *Nature Phys.* **12** (2015) 144-150.
17. ^{†*}Development of non-metallic diamond anvil cell and quantum oscillation measurement of CePtIn₇ in a pulsed-magnet: A. Miyake, Y. Kohama, S. Ohta, Y. Hirose, R. Settai, K. Matsubayashi, Y. Uwatoko, A. Matsuo, K. Kindo and M. Tokunaga, *J. Phys.: Conf. Ser.* **592** (2015) 012149(1-6).
18. ^{*}Synchrotron X-ray spectroscopy study on the valence state and magnetization in α -YbAl_{1-x}Fe_xB₄ ($x = 0.115$) at low temperatures and high magnetic fields: T. Terashima, Y. H. Matsuda, K. Kuga, S. Suzuki, Y. Matsumoto, S. Nakatsuji, A. Kondo, K. Kindo, N. Kawamura, M. Mizumaki and T. Inami, *J. Phys.: Conf. Ser.* **592** (2015) 012020 (6 pages).
19. ^{*}Field-induced quantum metal-insulator transition in the pyrochlore iridate Nd₂Ir₂O₇: Z. Tian, Y. Kohama, T. Tomita, H. Ishizuka, T. H. Hsieh, J. J. Ishikawa, K. Kindo, L. Balents and S. Nakatsuji, *Nature Phys.* **12** (2015) 134-139.
20. ^{†*}Magnetic and Structural Studies on Two-Dimensional Antiferromagnets (MCl)LaNb₂O₇ (M = Mn, Co, Cr): A. Kitada, Y. Tsujimoto, M. Nishi, A. Matsuo, K. Kindo, Y. Ueda, Y. Ajiro and H. Kageyama, *J. Phys. Soc. Jpn.* **85** (2016) 034005(6).
21. [†]Field-driven successive phase transitions in the quasi-two-dimensional frustrated antiferromagnet Ba₂CoTeO₆ and highly degenerate classical ground states: P. Chanlert, N. Kurita, H. Tanaka, D. Goto, A. Matsuo and K. Kindo, *Phys. Rev. B* **93** (2016) 094420(7).
22. ^{*}Experimental exploration of novel semimetal state in strong anisotropic Pyrochlore iridate Nd₂Ir₂O₇ under high magnetic field: Z. M. Tian, Y. Kohama, T. Tomita, J. Ishikawa, H. Mairo, K. Kindo and S. Nakatsuji, *J. Phys.: Conf. Ser.* **683** (2016) 012024(6).
23. ^{*}Frustrated magnetism in a Mott insulator based on a transition metal chalcogenide: S. Kawamoto, T. Higo, T. Tomita, S. Suzuki, Z. M. Tian, K. Mochizuki, A. Matsuo, K. Kindo and S. Nakatsuji, *J. Phys.: Conf. Ser.* **683** (2016) 012025(4).

Tokunaga group

We are focusing on physics in the quantum limit state, in which all the carriers are accommodated in the lowest Landau level. Since the carriers are confined in the smallest cyclotron orbit, kinetic degree of freedom normal to the field is suppressed. Thereby, the ratio between Coulomb interaction and bandwidth becomes large. Such strong correlation in the quantum limit state can realize anomalous quantum state. We can realize the quantum limit state in some kinds of semimetals with using non-destructive pulse magnets installed at ISSP. We studied magnetization and transport properties on various types of graphite in pulsed magnetic fields up to 74 T. The results suggest that the quantum limit state is realized in magnetic fields greater than 53 T applied along the *c*-axis. From systematic experimental studies, we proposed emergence of excitonic BCS-like state in

* Joint research among groups within ISSP.

the quantum limit state of graphite. We also studied transport properties of the semimetallic black phosphorus under pressure. We found anomalously large positive magnetoresistance and quantum oscillations in this material. The observed small Fermi surfaces, high mobilities, and light effective masses of carriers in semimetallic black phosphorus are comparable to those in the representative elemental semimetals of bismuth and graphite.

1. ^{†*}Anomalous Quantum Transport Properties in Semimetallic Black Phosphorus: K. Akiba, A. Miyake, Y. Akahama, K. Matsubayashi, Y. Uwatoko, H. Arai, Y. Fuseya and M. Tokunaga, *J. Phys. Soc. Jpn.* **84** (2015) 073708(1-4).
2. ^{*}Field Evolution of Quantum Critical and Heavy Fermi-Liquid Components in the Magnetization of the Mixed Valence Compound β -YbAlB₄: Y. Matsumoto, K. Kuga, Y. Karaki, Y. Shimura, T. Sakakibara, M. Tokunaga, K. Kindo and S. Nakatsuji, *J. Phys. Soc. Jpn.* **84** (2015) 024710(1-7).
3. ^{*}Possible Excitonic Phase of Graphite in the Quantum Limit State: K. Akiba, A. Miyake, H. Yaguchi, A. Matsuo, K. Kindo and M. Tokunaga, *J. Phys. Soc. Jpn.* **84** (2015) 054709(1-6).
4. High field studies on BiFeO₃ single crystals grown by the laser-diode heating floating zone method: M. Tokunaga, M. Akaki, A. Miyake, T. Ito and H. Kuwahara, *J. Magn. Magn. Mater.* **383** (2015) 259-261.
5. Magnetic field induced polar phase in the chiral magnet CsCuCl₃: A. Miyake, J. Shibuya, M. Akaki, H. Tanaka and M. Tokunaga, *Phys. Rev. B* **92** (2015) 100406(1-5).
6. ^{*}One-Third Magnetization Plateau with a Preceding Novel Phase in Volborthite: H. Ishikawa, M. Yoshida, K. Nawa, M. Jeong, S. Kramer, M. Horvatic, C. Berthier, M. Takigawa, M. Akaki, A. Miyake, M. Tokunaga, K. Kindo, J. Yamaura, Y. Okamoto and Z. Hiroi, *Phys. Rev. Lett.* **114** (2015) 227202(1-5).
7. ^{*}パルス強磁場下における磁気熱効果／比熱測定手法の開発：木原 工，小濱 芳允，徳永 将史，金道 浩一，*固体物理* **50** (2015) 371.
8. ハイスピードカメラで調べる強磁場物性：徳永 将史，*固体物理* **50** (2015) 163.
9. ^{†*}Superconductivity protected by spin–valley locking in ion-gated MoS₂: Y. Saito, Y. Nakamura, M. S. Bahrany, Y. Kohama, J. Ye, Y. Kasahara, Y. Nakagawa, M. Onga, M. Tokunaga, T. Nojima, Y. Yanase and Y. Iwasa, *Nature Phys.* **12** (2015) 144-150.
10. ^{†*}Development of non-metallic diamond anvil cell and quantum oscillation measurement of CePt₂In₇ in a pulsed-magnet: A. Miyake, Y. Kohama, S. Ohta, Y. Hirose, R. Settai, K. Matsubayashi, Y. Uwatoko, A. Matsuo, K. Kindo and M. Tokunaga, *J. Phys.: Conf. Ser.* **592** (2015) 012149(1-6).
11. Magnetic control of transverse electric polarization in BiFeO₃: M. Tokunaga, M. Akaki, T. Ito, S. Miyahara, A. Miyake, H. Kuwahara and N. Furukawa, *Nat. Commun.* **6** (2015) 5878(1-5).
12. [†]Thermodynamics and Kinetics of Martensitic Transformation in Ni-Mn-based Magnetic Shape Memory Alloys: X. Xu, R. Kainuma, T. Kihara, W. Ito, M. Tokunaga, T. Kanomata, N. Schryvers and J. Van Humbeeck, *MATEC Web of Conferences* **33** (2015) 01004(1-9).
13. [†]Spin Structure Change in Co-Substituted BiFeO₃: H. Yamamoto, T. Kihara, K. Oka, M. Tokunaga, K. Mibu and M. Azuma, *J. Phys. Soc. Jpn.* **85** (2016) 064704(1-4).
14. 圧力下の半金属黒磷における異常量子輸送現象：秋葉 和人，三宅 厚志，徳永 将史，赤浜 裕一，*固体物理* **51** (2016) 249.
15. Resistive memory effects in BiFeO₃ single crystals controlled by transverse electric fields: S. Kawachi, H. Kuroe, T. Ito, A. Miyake and M. Tokunaga, *Appl. Phys. Lett.* **108** (2016) 162903(1-4).
16. [†]Quantum Hall effect in a bulk antiferromagnet EuMnBi₂ with magnetically confined two-dimensional Dirac fermions: H. Masuda, H. Sakai, M. Tokunaga, Y. Yamasaki, A. Miyake, J. Shiogai, S. Nakamura, S. Awaji, A. Tsukazaki, H. Nakao, Y. Murakami, T. -h. Arima, Y. Tokura and S. Ishiwata, *Science Advances* **2** (2016) e1501117(1-6).

Y. Matsuda group

The valence state and magnetization of the heavy fermion compound α -YbAlB₄ have been investigated. This compound shows the strong valence fluctuation and its Yb valence is around 2.8 which is distinctly smaller than the expected valence state 3.0 for local magnetic Yb ions. Although the fluctuated valence is expected to be influenced by magnetic field due to the suppression of the Kondo effect, the valence state has been found to be almost independent of magnetic field even at rather high magnetic field of up to 40 T. This experimental fact contrasts to that obtained in a related antiferromagnetic compound α -YbAl_{1-x}Fe_xB₄ ($x = 0.115$); the small valence increase is observed when magnetic field exceeds 20 T. The quantum criticality might be

[†] Joint research with outside partners.

enhanced in YbAlB₄ at low temperatures and the criticality is not significant in Fe-doped substance, which may explain the different magnetic field dependence of the Yb valence. In addition to the study of YbAlB₄, the B-T phase diagram of solid oxygen has been clarified and the high-field θ -phase has found to have low entropy and be clearly different from the high-temperature γ -phase. We also studied insulator-metal transition of Co-oxides, Fe-based superconductors and magneto-optics of the multiferroic CuB₂O₄.

1. *Magnetic-Field-Induced Insulator–Metal Transition in (Pr_{1-y}Y_y)_{0.7}Ca_{0.3}CoO₃ at Ultrahigh Magnetic Fields: S. Lee, Y. H. Matsuda, T. Naito, D. Nakamura and S. Takeyama, *J. Phys. Soc. Jpn.* **84** (2015) 044703-7.
2. *X-ray Absorption Spectroscopy in the Heavy Fermion Compound α -YbAlB₄ at High Magnetic Fields: T. T. Terashima, Y. H. Matsuda, K. Kuga, S. Suzuki, Y. Matsumoto, S. Nakatsuji, A. Kondo, K. Kindo, N. Kawamura, M. Mizumaki and T. Inami, *J. Phys. Soc. Jpn.* **84** (2015) 114715(1-4).
3. †*Phase boundary of θ phase of solid oxygen in ultrahigh magnetic fields: T. Nomura, Y. H. Matsuda, S. Takeyama, A. Matsuo, K. Kindo and T. C. Kobayashi, *Phys. Rev. B* **92** (2015) 064109 (4 pages).
4. †*One-Way Transparency of Light in Multiferroic CuB₂O₄: S. Toyoda, N. Abe, S. Kimura, Y. H. Matsuda, T. Nomura, A. Ikeda, S. Takeyama and T. Arima, *Phys. Rev. Lett.* **115** (2015) 267207 (5 pages).
5. †*Anisotropy in the upper critical field of FeSe and FeSe_{0.33}Te_{0.67} single crystals: J. L. Her, Y. Kohama, Y. H. Matsuda, K. Kindo, W.-H. Yang, D. A. Chareev, E. S. Mitrofanova, O. S. Volkova, A. N. Vasiliev and J.-Y. Lin, *Supercond. Sci. Technol.* **28** (2015) 045013 (6 pages).
6. 放射光 X 線実験で磁場を使う：松田 康弘，*放射光* **28** (2015) 221-228.
7. † 超強磁場で誘起される固体酸素の新しい相：松田 康弘，*パリティ* **30** (2015) 48-51.
8. †*Anomalous diamagnetic shifts in InP-GaP lateral quantum-wires: Y. H. Shin, B. K. Choi, Y. Kim, J. D. Song, D. Nakamura, Y. H. Matsuda and S. Takeyama, *Opt. Express* **23** (2015) 28349-28357.
9. *Synchrotron X-ray spectroscopy study on the valence state and magnetization in α -YbAl_{1-x}Fe_xB₄ ($x = 0.115$) at low temperatures and high magnetic fields: T. Terashima, Y. H. Matsuda, K. Kuga, S. Suzuki, Y. Matsumoto, S. Nakatsuji, A. Kondo, K. Kindo, N. Kawamura, M. Mizumaki and T. Inami, *J. Phys.: Conf. Ser.* **592** (2015) 012020 (6 pages).
10. †Probe of the Band Structure of MBE Grown p-Type InMnAs at Ultrahigh Magnetic Fields: Y. Sun, F. V. Kyrychenko, G. D. Sanders, C. J. Stanton, G. A. Khodaparast, J. Kono, Y. H. Matsuda and H. Munekata, *SPIN* **05** (2015) 1550002 (16 pages).

Center of Computational Materials Science

Akai group

(1) It is crucial to treat f-states properly to describe the magnetic properties of rare-earth permanent magnet materials. However, the local or semi-local approximations to the density functional method, which is the common basis of first-principles electronic structure calculations, are unable to treat the f-states in a reasonable way. To overcome this situation we have developed several methods that go beyond the local density approximation. One of them are the optimized effective potential method combined with the exact-exchange and random-phase approximation. Another is the self-interaction corrected LDA/GGA applied to f-states. Using these methods, the magnetic crystalline anisotropy of Sm₂Fe₁₇N_x ($0 < x < 3$) is calculated. The results obtained by two different methods show considerable differences and the origin of these discrepancies are discussed. (2) A scheme that combines the non-equilibrium Green's function method with the Korringa-Kohn-Rostoker (KKR) Green's function method is proposed. The scheme applied to Schottky contact composed of Al/GaN/Al trilayer. The transport property of this system under various finite bias voltages is calculated. It is shown that the asymmetric behavior of electron transport against the direction of bias voltage occurs in this system, confirming the feature of rectification.

1. Ab initio study of ⁵⁹Co NMR spectra in Co₂FeAl_{1-x}Si_x Heusler alloys: H. Nishihara, K. Sato, H. Akai, C. Takiguchi, M. Geshi, T. Kanomata, T. Sakon and T. Wada, *Physica B* **485** (2015) 66-70.
2. Near-field correction in the first-principles calculations by the exact two-center expansion for the inverse of the distance: M. Ogura, C. Zecha, M. Offenberger, H. Ebert and H. Akai, *J. Phys.: Condens. Matter* **27** (2015) 485201(1-8).
3. Optimized effective potential method and application to a static RPA correlation: T. Fukazawa and H. Akai, *J. Phys.: Condens. Matter* **27** (2015) 115502(1-10).

* Joint research among groups within ISSP.

4. Formulation of the augmented plane-wave and muffin-tin orbital method: T. Kotani, H. Kino and H. Akai, *J. Phys. Soc. Jpn.* **84** (2015) 034701(1-9).
5. Role of N in the Permanent Magnet Material $\text{Sm}_2\text{Fe}_{17}\text{N}_x$: M. Ogura, A. Mashiyama and H. Akai, *J. Phys. Soc. Jpn.* **84** (2015) 084702(1-6).
6. The metamagnetic behavior and giant inverse magnetocaloric effect in Ni–Co–Mn–(Ga, In, Sn) Heusler alloys: P. Entel, V. V. Sokolovskiy, V. D. Buchelnikov, M. Ogura, M. E. Gruner, A. Gruenebohm, D. Comtesse and H. Akai, *J. Magn. Mater.* **385** (2015) 193-197.
7. Relevance of 4f-3d exchange to finite-temperature magnetism of rare-earth permanent magnets: An ab-initio-based spin model approach for NdFe_{12}N : M. Matsumoto, H. Akai, Y. Harashima, S. Doi and T. Miyake, *J. Appl. Phys.* **119** (2016) 213901(1-7).

Ozaki Taisuke group

First-principles calculations based on density functional theories (DFT) have been playing an invaluable role as a cornerstone in computational materials science. For a wide variety of materials ranging from metals, insulators, semiconductors, and molecules the DFT calculations enable us to quantitatively predict the chemical and physical properties within a reasonable computational cost. With a recent advance of massively parallel computers, even realistic materials discussed in industrial fields have been becoming the potential targets. However, it is a challenging problem to develop algorithms and software being suitable for massively parallel computers typified by the K-computer, which results in a fact that users and developers are distinguished from each other in recent years, and that most of users use specific software. We have been developing OpenMX (Open source package for Material eXplore) towards a de fact standard DFT code, and in 2015 released the Ver. 3.8 to the public under GNU-GPL. To improve numerical accuracy, optimized pseudopotentials and pseudo-atomic basis functions have been developed based on a norm-conserving pseudopotential method with multiple reference energies and a variational optimization method for radial basis functions. The optimization over 80 elements in the periodic table results in the reliable database storing optimized pseudopotentials and pseudo-atomic basis functions, where the accuracy of the database was validated by the delta gauge method. In addition, new functionalities including stress tensor calculation, band unfolding method, and eigenchannel/real space current analysis have been developed, which makes OpenMX a versatile tool for many applications.

1. First-Principles Study on Cubic Pyrochlore Iridates $\text{Y}_2\text{Ir}_2\text{O}_7$ and $\text{Pr}_2\text{Ir}_2\text{O}_7$: F. Ishii, Y. P. Mizuta, T. Kato, T. Ozaki, H. Weng and S. Onoda, *J. Phys. Soc. Jpn.* **84** (2015) 073703(5 pages).
2. Reproducibility in density functional theory calculations of solids: K. Lejaeghere, G. Bihlmayer, T. Bjorkman, P. Blaha, S. Blugel, V. Blum, D. Caliste, I. E. Castelli, S. J. Clark, A. Dal Corso, S. de Gironcoli, T. Deutsch, J. K. Dewhurst, I. Di Marco, C. Draxl, M. Du ak, O. Eriksson, J. A. Flores-Livas, K. F. Garrity, L. Genovese, P. Giannozzi, M. Giantomassi, S. Goedecker, X. Gonze, O. Granas, E. K. U. Gross, A. Gulans, F. Gygi, D. R. Hamann, P. J. Hasnip, N. A. W. Holzwarth, D. Iu an, D. B. Jochym, F. Jollet, D. Jones, G. Kresse, K. Koepnik, E. Kucukbenli, Y. O. Kvashnin, I. L. M. Locht, S. Lubeck, M. Marsman, N. Marzari, U. Nitzsche, L. Nordstrom, T. Ozaki, L. Paulatto, C. J. Pickard, W. Poelmans, M. I. J. Probert, K. Refson, M. Richter, G. -M. Rignanese, S. Saha, M. Scheffler, M. Schlipf, K. Schwarz, S. Sharma, F. Tavazza, P. Thunstrom, A. Tkatchenko, M. Torrent, D. Vanderbilt, M. J. van Setten, V. Van Speybroeck, J. M. Wills, J. R. Yates, G. -X. Zhang and S. Cottenier, *Science* **351** (2016) aad3000(1-7).
3. Hybrid and 4-D FFT implementations of an open-source parallel FFT package OpenFFT: T. V. T. Duy and T. Ozaki, *J. Supercomput* **72** (2016) 391-416.

Laser and Synchrotron Research Center

Suemoto group

(1) We tried to control the orientation of macroscopic magnetization in an orthoferrite ErFeO_3 by using terahertz magnetic field enhanced by a split ring resonator. The magnetization orientation was successfully controlled by changing the time delay between the terahertz pulse and the heating pulse. In addition, it was found that two kinds of mechanisms, i.e., spin precession motion and oscillating external magnetic field, are relevant on the symmetry breaking at the beginning of spin reorientation phase transition induced by the heating pulse. This is probably the first demonstration of coherent control of the macroscopic magnetization using direct magnetic dipole interaction with radiation. (2) Femtosecond infrared luminescence was observed in a topological insulator (TlBiSe_2) and the luminescence component below the band gap energy (0.35 eV) was ascribed to the carrier recombination in the metal-like two-dimensional Dirac bands at the surface, while that above 0.35eV was assigned to the transitions in the semiconductor-like bulk bands. These results show that the luminescence method is usable under ambient condition for investigation of carrier dynamics, while photoemission method requires ultrahigh vacuum, which is different from

[†] Joint research with outside partners.

the operand condition of realistic devices.

1. Dynamics of photoinduced change of magneto-anisotropy parameter in orthoferrites probed with terahertz excited coherent spin precession: K. Yamaguchi, T. Kurihara, H. Watanabe, M. Nakajima and T. Suemoto, *Phys. Rev. B* **92** (2015) 064404.
2. Magnetization-free measurements of spin orientations in orthoferrites using terahertz time domain spectroscopy: T. Suemoto, K. Nakamura, T. Kurihara and H. Watanabe, *Appl. Phys. Lett.* **107** (2015) 042404.
3. [†]Ultraviolet stimulated emission from high-temperature-annealed MgO microcrystals at room temperature: H. Soma, Y. Uenaka, A. Asahara, T. Suemoto and T. Uchino, *Appl. Phys. Lett.* **106** (2015) 041116(1-5).
4. ^{*}Transient gain analysis of gain-switched semiconductor lasers during pulse lasing: S. Chen, T. Ito, A. Asahara, H. Nakamae, T. Nakamura, M. Yoshita, C. Kim, B. Zhang, H. Yokoyama, T. Suemoto and H. Akiyama, *Appl. Opt.* **54** (2015) 10438.
5. ^{†*}Layer number dependence of carrier lifetime in graphenes observed using time-resolved mid-infrared luminescence: H. Watanabe, T. Kawasaki, T. Iimori, F. Komori and T. Suemoto, *Chem. Phys. Lett.* **637** (2015) 58-62.
6. ^{*}Optically detecting the edge-state of a three-dimensional topological insulator under ambient conditions by ultrafast infrared photoluminescence spectroscopy: S.-Y. Maezawa, H. Watanabe, M. Takeda, K. Kuroda, T. Someya, I. Matsuda and T. Suemoto, *Sci. Rep.* **5** (2015) 16443(1-8).
7. [†]Investigation of insulator-metal transition in Ti₄O₇ using terahertz probe pulse: H. Kamioka, J. Nishitani, H. Tsukada, R. Yamaguchi and T. Suemoto, *Appl. Phys. Lett.* **108** (2016) 071908.
8. ^{*}Optical pump-THz probe analysis of long-lived *d*-electrons and relaxation to self-trapped exciton states in MnO: J. Nishitani, T. Nagashima, M. Lippmaa and T. Suemoto, *Appl. Phys. Lett.* **108** (2016) 162101(1-5).
9. ^{*}Photoexcited *d*-electron dynamics in transition metal oxide MnO studied by optical pump-THz probe measurements: J. Nishitani, T. Kurihara, A. Asahara, T. Nagashima, M. Lippmaa and T. Suemoto, *Phys. Status Solidi C* **13** (2016) 113-116.
10. [†]Magnetic-excitation-assisted photoluminescence from self-trapped exciton states in MnO: J. Nishitani, T. Nagashima and T. Suemoto, *J. Phys. Condens. Matter* **28** (2016) 016004.
11. [†]Dynamics of photoinduced phase transitions in a Prussian blue analog studied by CN vibrational spectroscopy (Review paper): T. Suemoto, R. Fukaya, A. Asahara, H. Watanabe, H. Tokoro and S.-I. Ohkoshi, *Current Inorganic Chemistry* **6** (2016) 10-25.

Shin group

We studied high T_c Fe-pnictide superconductors using 7-eV laser. High resolution photoemission study with polarization dependence is very powerful for the study of the superconducting mechanism. Orbital fluctuation mechanism is also important in addition to the spin fluctuation mechanism.

1. ^{†*}Low-Temperature and High-Energy-Resolution Laser Photoemission Spectroscopy: T. Shimojima, K. Okazaki and S. Shin, *J. Phys. Soc. Jpn.* **84** (2015) 072001(1-26).
2. [†]Hybridization gap formation in the Kondo insulator YbB₁₂ observed using time-resolved photoemission spectroscopy: M. Okawa, Y. Ishida, M. Takahashi, T. Shimada, F. Iga, T. Takabatake, T. Saitoh and S. Shin, *Phys. Rev. B* **92** (2015) 161108(1-5).
3. ^{†*}Nonequilibrium electronic and phonon dynamics of Cu_{0.17}Bi₂Se₃ investigated by core-level and valence-band time-resolved photoemission spectroscopy: T. Yamamoto, Y. Ishida, R. Yoshida, M. Okawa, K. Okazaki, T. Kanai, A. Kikkawa, Y. Taguchi, T. Kiss, K. Ishizaka, N. Ishii, J. Itatani, S. Watanabe, Y. Tokura, T. Saitoh and S. Shin, *Phys. Rev. B* **92** (2015) 121106(1-6).
4. ^{*}Ultrafast spin-switching of a ferrimagnetic alloy at room temperature traced by resonant magneto-optical Kerr effect using a seeded free electron laser: Sh. Yamamoto, M. Taguchi, T. Someya, Y. Kubota, S. Ito, H. Wadati, M. Fujisawa, F. Capotondi, E. Pedersoli, M. Manfredda, L. Raimondi, M. Kiskinova, J. Fujii, P. Moras, T. Tsuyama, T. Nakamura, T. Kato, T. Higashide, S. Iwata, S. Yamamoto, S. Shin and I. Matsuda, *Rev. Sci. Instrum.* **86** (2015) 083901(1-5).
5. [†]Ultrahigh-spatial-resolution chemical and magnetic imaging by laser-based photoemission electron microscopy: T. taniuchi, Y. Kotani and S. Shin, *Rev. Sci. Instrum.* **86** (2015) 023701(5 pages).

* Joint research among groups within ISSP.

6. † Gigantic Surface Lifetime of an Intrinsic Topological Insulator: M. Neupane, S.-Y. Xu, Y. Ishida, S. Jia, B. M. Fregoso, C. Liu, I. Belopolski, G. Bian, N. Alidoust, T. Durakiewicz, V. Galitski, S. Shin, R. J. Cava and M. Zahid Hasan, *Phys. Rev. Lett.* **115** (2015) 116801(1-5).
7. † Temperature Dependence of Magnetically Active Charge Excitations in Magnetite across the Verwey Transition: M. Taguchi, A. Chainani, S. Ueda, M. Matsunami, Y. Ishida, R. Eguchi, S. Tsuda, Y. Takata, M. Yabashi, K. Tamasaku, Y. Nishino, T. Ishikawa, H. Daimon, S. Todo, H. Tanaka, M. Oura, Y. Senba, H. Ohashi and S. Shin, *Phys. Rev. Lett.* **115** (2015) 256405(1-5).
8. †* フェムト秒域時間分解光電子分光法によるグラフェンの超高速キャリアダイナミクスの追跡 Tracing Ultrafast Carrier Dynamics in Graphene with Femtosecond Time-resolved Photoemission Spectroscopy: 染谷 隆史, 吹留 博一, 石田 行章, 吉田 力矢, 山本 達, 板谷 治郎, 小森 文夫, 辛 埴, 松田 巖, *表面科学* **36(8)** (2015) 418-423.
9. † Reaction of Sb on In/Si(111) surfaces: Heteroepitaxial InSb(111) formation: M. Hashimoto, A. Nakaguchi, F. -Z. Guo, M. Ueda, T. Yasue, T. Matsushita, T. Kinoshita, K. Kobayashi, M. Oura, T. Takeuchi, Y. Saito, S. Shin and T. Koshikawa, *Surf. Sci.* **641** (2015) 121-127.
10. †* 高次高調波の固体時間分解光電子分光への応用 Application of High Harmonic Generation to Time-Resolved Photoemission Spectroscopy of Solids: 岡崎 浩三, 辛 埴, *レーザー研究* **43 (12)** (2015) 838-843.
11. †* Selective Formation of Zigzag Edges in Graphene Cracks: M. Fujihara, R. Inoue, R. Kurita, T. Taniuchi, Y. Motoyui, S. Shin, F. Komori, Y. Maniwa, H. Shinohara and Y. Miyata, *ACS Nano* **9** (2015) 9027-9033.
12. †* Point nodes persisting far beyond T_c in Bi2212: T. Kondo, W. Malaeb, Y. Ishida, T. Sasagawa, H. Sakamoto, T. Takeuchi, T. Tohyama and S. Shin, *Nat. Commun.* **6** (2015) 7699(8 pages).
13. †* Quadratic Fermi Node in a 3D Strongly Correlated Semimetal: T. Kondo, M. Nakayama, R. Chen, J. J. Ishikawa, E. -G. Moon, T. Yamamoto, Y. Ota, W. Malaeb, H. Kanai, Y. Nakashima, Y. Ishida, R. Yoshida, H. Yamamoto, M. Matsunami, S. Kimura, N. Inami, K. Ono, H. Kumigashira, S. Nakatsuji, L. Balents and S. Shin, *Nat. Commun.* **6** (2015) 10042(1-8).
14. †* Emergent photovoltage on SmB_6 surface upon bulk-gap evolution revealed by pump-and-probe photoemission spectroscopy: Y. Ishida, T. Otsu, T. Shimada, M. Okawa, Y. Kobayashi, F. Iga, Y. Takabatake and S. Shin, *Sci. Rep.* **5** (2015) 8160(1-6).
15. †* Ultrafast electron dynamics at the Dirac node of the topological insulator Sb_2Te_3 : S. Zhu, Y. Ishida, K. Kuroda, K. Sumida, M. Ye, J. Wang, H. Pan, M. Taniguchi, S. Qiao, S. Shin and A. Kimura, *Sci. Rep.* **5** (2015) 13213(1-6).
16. † Electronic Structure Evolution across the Peierls Metal-Insulator Transition in a Correlated Ferromagnet: P. A. Bhobe, A. Kumar, M. Taguchi, R. Eguchi, M. Matsunami, Y. Takata, A. K. Nandy, P. Mahadevan, D. D. Sarma, A. Neroni, E. Sasioglu, M. Lezaic, M. Oura, Y. Senba, H. Ohashi, K. Ishizaka, M. Okawa, S. Shin, K. Tamasaku, Y. Kohmura, M. Yabashi, T. Ishikawa, K. Hasegawa, M. Isobe, Y. Ueda and A. Chainani, *Phys. Rev. X* **5** (2015) 041004(1-9).
17. †* One-dimensional metallic surface states of Pt-induced atomic nanowires on Ge(001): K. Yaji, S. Kim, I. Mochizuki, Y. Takeichi, Y. Ohtsubo, P. L. Fèvre, F. Bertran, A. Taleb-Ibrahimi, S. Shin and F. Komori, *J. Phys.: Condens. Matter* **28** (2016) 284001(1-9).
18. † Coexistence of a pseudo-gap and a superconducting gap for the high- T_c superconductor LSCO using photoemission spectroscopy: T. Yoshida, W. Malaeb, S. Ideta, D. H. Lu, R. G. Moor, Z. -X. Shen, M. Okawa, T. Kiss, K. Ishizaka, S. Shin, S. Komiya, Y. Ando, H. Eisaki, S. Uchida and A. Fujimori, *Phys. Rev. B* **93** (2016) 014513(5 pages).
19. † Revealing the ultrafast light-to-matter energy conversion before heat diffusion in a layered Dirac semimetal: Y. Ishida, H. Masuda, H. Sakai, S. Ishiwata and S. Shin, *Phys. Rev. B* **93** (2016) 100302(6 pages).
20. * High-resolution three-dimensional spin- and angle-resolved photoelectron spectrometer using vacuum ultraviolet laser light: K. Yaji, A. Harasawa, K. Kuroda, S. Toyohisa, M. Nakayama, Y. Ishida, A. Fukushima, S. Watanabe, C. Chen, F. Komori and S. Shin, *Rev. Sci. Instrum.* **87** (2016) 053111(1-6).
21. †* Spin Polarization and Texture of the Fermi Arcs in the Weyl Fermion Semimetal TaAs: S.-Y. Xu, I. Belopolski, D. S. Sanchez, M. Neupane, G. Chang, K. Yaji, Z. Yuan, C. Zhang, K. Kuroda, G. Bian, C. Guo, H. Lu, T.-R. Chang, N. Alidoust, H. Zheng, C.-C. Lee, S.-M. Huang, C.-H. Hsu, H.-T. Jeng, A. Bansil, T. Neupert, F. Komori, T. Kondo, S. Shin, H. Lin, S. Jia and M. Zahid Hasan, *Phys. Rev. Lett.* **116** (2016) 096801(1-7).
22. †* 角度分解光電子分光による精密測定で解き明かす銅酸化物高温超伝導体の擬ギャップと超伝導ギャップの競合関係: 近藤 猛, 竹内 恒博, 辛 埴, *固体物理* **51** (2016) 203-221.
23. 超高速時間分解光電子分光: 石田 行章, *表面科学* **37** (2016) 31-36.

† Joint research with outside partners.

24. [†]Electronic structure and relaxation dynamics in a superconducting topological material: M. Neupane, Y. Ishida, R. Sankar, J.-X. Zhu, D. S. Sanchez, I. Belopolski, S.-Y. Xu, N. Alidoust, M. Mofazzel Hosen, S. Shin, F. Chou, M. Zahid Hasan and T. Durakiewicz, *Sci. Rep.* **6** (2016) 22557(7 pages).
25. [†]Quasi-particles ultrafastly releasing kink bosons to form Fermi arcs in a cuprate superconductor: Y. Ishida, T. Saitoh, T. Mochiku, T. Nakane, K. Hirata and S. Shin, *Sci. Rep.* **6** (2016) 18747(8 pages).

Akiyama group

In 2015, we studied radiation damage effects of in multi-junction solar cells via absolute electroluminescence-efficiency measurements, by developing LED radiance secondary standards. We studied pico- and femto-second short-pulse generation via gain switching in GaAs double-hetero semiconductor lasers. We studied quantum yields of equarin in jelly-fish bioluminescence. We also made intensive studies on theoretical quantum-chemistry and molecular-dynamics calculations on oxyluciferins.

1. Time-resolved observation of coherent excitonic nonlinear response with a table-top narrowband THz pulse wave: K. Uchida, H. Hirori, T. Aoki, C. Wolpert, T. Tamaya, K. Tanaka, T. Mochizuki, C. Kim, M. Yoshita, H. Akiyama, L. N. Pfeiffer and K. W. West, *Appl. Phys. Lett.* **107** (2015) 221106.
2. Time-resolved photoluminescence measurements for determining voltage-dependent charge-separation efficiencies of subcells in triple-junction solar cells: D. M. Tex, T. Ihara, H. Akiyama, M. Imaizumi and Y. Kanemitsu, *Appl. Phys. Lett.* **106** (2015) 013905.
3. 多接合太陽電池の診断・設計と発光絶対値・発光量子効率の評価：秋山 英文，*応用物理* **84** (2015) 319-325.
4. ^{*}Transient gain analysis of gain-switched semiconductor lasers during pulse lasing: S. Chen, T. Ito, A. Asahara, H. Nakamae, T. Nakamura, M. Yoshita, C. Kim, B. Zhang, H. Yokoyama, T. Suemoto and H. Akiyama, *Appl. Opt.* **54** (2015) 10438.
5. ^{*}Effect of very high magnetic field on the optical properties of firefly light emitter oxyluciferin: W. Zhou, D. Nakamura, Y. Wang, T. Mochizuki, H. Akiyama and S. Takeyama, *J. Lumin.* **165** (2015) 15.
6. Multi-junction-solar-cell designs and characterizations based on detailed-balance principle and luminescence yields: H. Akiyama, L. Zhu, M. Yoshita, C. Kim, S. Chen, T. Mochizuki and Y. Kanemitsu, *Proc. SPIE* **9358** (2015) 93580B1-8.
7. Time-resolved observation of excitonic dynamics under coherent terahertz excitation in GaAs quantum wells: K. Uchida, H. Hirori, T. Aoki, C. Wolpert, K. Tanaka, T. Mochizuki, C. Kim, M. Yoshita, H. Akiyama, L. N. Pfeiffer and K. W. West, *Proc. SPIE* **9361** (2015) 93611L1-6.
8. Analysis of Oxyluciferin Photoluminescence Pathways in Aqueous Solutions: M. Hiyama, T. Mochizuki, H. Akiyama and N. Koga, *Photochem. Photobiol.* **91** (2015) 74.
9. ^{*}Vibronic Structures in Absorption and Fluorescence Spectra of Firefly Oxyluciferin in Aqueous Solutions: M. Hiyama, Y. Noguchi, H. Akiyama, K. Yamada and N. Koga, *Photochem. Photobiol.* **91** (2015) 819.
10. Degradation mechanism of perovskite $\text{CH}_3\text{NH}_3\text{PbI}_3$ diode devices studied by electroluminescence and photoluminescence imaging spectroscopy: M. Okano, M. Endo, A. Wakamiya, M. Yoshita, H. Akiyama and Y. Kanemitsu, *Appl. Phys. Express* **8** (2015) 102302.
11. On the importance of cavity-length and heat dissipation in GaN-based vertical-cavity surface-emitting lasers: W. J. Liu, X. L. Hu, L. Y. Ying, S. Q. Chen, J. Y. Zhang, H. Akiyama, Z. P. Cai and B. P. Zhang, *Sci. Rep.* **5** (2015) 9600.
12. Thorough subcells diagnosis in a multi-junction solar cell via absolute electroluminescence-efficiency measurements: S. Chen, L. Zhu, M. Yoshita, T. Mochizuki, C. Kim, H. Akiyama, M. Imaizumi and Y. Kanemitsu, *Sci. Rep.* **5** (2015) 7836.
13. ^{*}First-Principles Investigation of Strong Excitonic Effects in Oxygen 1s X-ray Absorption Spectra: Y. Noguchi, M. Hiyama, H. Akiyama, Y. Harada and N. Koga, *J. Chem. Theory Comput.* **11** (2015) 1668-1673.
14. Strong localization effect and carrier relaxation dynamics in self-assembled InGaN quantum dots emitting in the green: G.-E. Weng, W.-R. Zhao, S.-Q. Chen, H. Akiyama, Z.-C. Li, J.-P. Liu and B.-P. Zhang, *Nanoscale Res. Lett.* **10** (2015) 31.
15. Absolute Electroluminescence Imaging of Multi-Junction Solar Cells and Calibration Standards: M. Yoshita, L. Zhu, C. Kim, H. Akiyama, S. Chen, T. Mochizuki, H. Kubota, T. Nakamura, M. Imaizumi and Y. Kanemitsu, *The 42nd IEEE Photovoltaic Specialists Conference Proceedings* **7356199** (2015) 1-4.

^{*} Joint research among groups within ISSP.

16. Characterizations of radiation damages in multi-junction solar cells focused on subcell internal luminescence quantum yields via absolute electroluminescence measurements: L. Zhu, M. Yoshita, S. Chen, T. Nakamura, T. Mochizuki, C. Kim, M. Imaizumi, Y. Kanemitsu and H. Akiyama, The 42nd IEEE Photovoltaic Specialists Conference Proceedings **7355671** (2015) 1-4.
17. Determining Subcell Carrier-Collection Efficiencies of Triple-Junction Solar Cells Using Time-Resolved Photoluminescence: D. M. Tex, T. Ihara, H. Akiyama, M. Imaizumi and Y. Kanemitsu, The 42nd IEEE Photovoltaic Specialists Conference Proceedings **7356197** (2015) 1-4.
18. Solar-cell radiance standard for absolute electroluminescence measurements and open-circuit voltage mapping of silicon solar modules: T. Mochizuki, C. Kim, M. Yoshita, J. Mitchell, Z. Lin, S. Chen, H. Takato, Y. Kanemitsu and H. Akiyama, J. Appl. Phys. **119** (2016) 034501.
19. Conversion efficiency limits and bandgap designs for multi-junction solar cells with internal radiative efficiencies below unity: L. Zhu, T. Mochizuki, M. Yoshita, S. Chen, C. Kim, H. Akiyama and Y. Kanemitsu, Opt. Express **24** (2016) A740-A751.
20. * High-precision group-delay dispersion measurements of optical fibers via fingerprint-spectral wavelength-to-time mapping: T. Ito, O. Slezak, M. Yoshita, H. Akiyama and Y. Kobayashi, Photon. Res. **4** (2016) 13-16.
21. Calibration standards and measurement accuracy of absolute electroluminescence and internal properties in multi-junction and arrayed solar cells: M. Yoshita, L. Zhu, C. Kim, T. Mochizuki, T. Nakamura, M. Imaizumi, S. Chen, H. Kubota, Y. Kanemitsu and H. Akiyama, in Proc. of SPIE Photonics West **9743** (2016) 97430D1-6.
22. Characterization and modeling of radiation damages via internal radiative efficiency in multi-junction solar cells: L. Zhu, M. Yoshita, T. Nakamura, M. Imaizumi, C. Kim, T. Mochizuki, S. Chen, Y. Kanemitsu and H. Akiyama, in Proc. of SPIE Photonics West **9743** (2016) 97430U1-7.
23. Characterizations of Radiation Damage in Multijunction Solar Cells Focused on Subcell Internal Luminescence Quantum Yields via Absolute Electroluminescence Measurements: L. Zhu, M. Yoshita, S. Chen, T. Nakamura, T. Mochizuki, C. Kim, M. Imaizumi, Y. Kanemitsu and H. Akiyama, IEEE J. Photovoltaics **6** (2016) 777-782.

I. Matsuda group

Developments and experiments of the advanced spectroscopies have been carried out by using vacuum ultraviolet (VUV) and soft X-rays (SX). At SPring-8 BL07LSU, we have succeeded in realizing fast-switching (10 Hz) of the light polarizations for the segmented cross-type undulator. Moreover, we could measure a spectrum of X-ray magnetic circular dichroism for a magnetic sample using the fast-switching of left and right circular polarized light. At the end-station, we routinely supported picosecond-time-resolved SX photoemission spectroscopy experiments of joint-researches. Studies of photovoltaics and photocatalysis have been carried out mainly. In the laboratory, we studied electronic structure of novel two-dimensional materials that could show intriguing dynamical properties.

1. Deeper insight into phase relations in ultrathin Pb films: R.-Y. Liu, A. Huang, C.-C. Huang, C.-Y. Lee, C.-H. Lin, C.-M. Cheng, K.-D. Tsuei, H.-T. Jeng, I. Matsuda and S. -J. Tang, Phys. Rev. B **92** (2015) 115415(1-8).
2. * Scanning tunneling spectroscopy study of quasiparticle interference on dual topological insulator $\text{Bi}_{1-x}\text{Sb}_x$: S. Yoshizawa, F. Nakamura, A. A. Taskin, T. Iimori, K. Nakatsuji, I. Matsuda, Y. Ando and F. Komori, Phys. Rev. B **91** (2015) 045423(1-6).
3. * Ultrafast spin-switching of a ferrimagnetic alloy at room temperature traced by resonant magneto-optical Kerr effect using a seeded free electron laser: Sh. Yamamoto, M. Taguchi, T. Someya, Y. Kubota, S. Ito, H. Wadati, M. Fujisawa, F. Capotondi, E. Pedersoli, M. Manfredda, L. Raimondi, M. Kiskinova, J. Fujii, P. Moras, T. Tsuyama, T. Nakamura, T. Kato, T. Higashide, S. Iwata, S. Yamamoto, S. Shin and I. Matsuda, Rev. Sci. Instrum. **86** (2015) 083901(1-5).
4. Photoemission Circular Dichroism and Spin Polarization of the Topological Surface States in Ultrathin Bi_2Te_3 Films: C. -Z. Xu, Y. Liu, R. Yukawa, L. -X. Zhang, I. Matsuda, T. Miller and T. -C. Chiang, Phys. Rev. Lett. **115** (2015) 016801 (1-5).
5. †* フェムト秒域時間分解光電子分光法によるグラフェンの超高速キャリアダイナミクスの追跡 Tracing Ultrafast Carrier Dynamics in Graphene with Femtosecond Time-resolved Photoemission Spectroscopy: 染谷 隆史, 吹留 博一, 石田 行章, 吉田 力矢, 山本 達, 板谷 治郎, 小森 文夫, 辛 埴, 松田 巖, 表面科学 **36(8)** (2015) 418-423.
6. Significantly enhanced giant Rashba splitting in a thin film of binary alloy: W.-C. Chen, T.-R. Chang, S.-T. Tai, S. Yamamoto, J.-M. Kuo, C.-M. Cheng, K.-D. Tsuei, K. Yaji, H. Lin, H.-T. Jeng, C.-Y. Mou, I. Matsuda and S.-J. Tang, New J. Phys. **17** (2015) 083015(1-12).

† Joint research with outside partners.

7. † グラフェンの超高速状態を直接観測：染谷 隆史，吹留 博一，松田 巖，光アイアンス **26** (2015) 31.
8. * Optically detecting the edge-state of a three-dimensional topological insulator under ambient conditions by ultrafast infrared photoluminescence spectroscopy: S.-Y. Maezawa, H. Watanabe, M. Takeda, K. Kuroda, T. Someya, I. Matsuda and T. Suemoto, *Sci. Rep.* **5** (2015) 16443 (1-8).
9. Microscopic observation and chemical mapping of opal phytoliths in a mulberry leaf: O. Tsutsui, R. Sakamoto, M. Hattori, K. Hasegawa, T. Handa, D. Nishio-Hamane and I. Matsuda, *Flora* **218** (2015) 44-50.
10. Anisotropic effective mass approach for multiple subband structures at wide-gap semiconductor surfaces: Application to accumulation layers of SrTiO₃ and ZnO: R. Yukawa, K. Ozawa, S. Yamamoto, R. -Y. Liu and I. Matsuda, *Surface Science* **641** (2015) 224-230.
11. Generation of metallic e_g -derived band at Cs/SrTiO₃ interface observed by polarization-dependent photoemission spectroscopy: K. Akikubo, I. Matsuda, D. Schmaus, G. Marcaud, S. Yamamoto, R.-Y. Liu, R. Yukawa, M. G. Silly, F. Sirotti and M. D'Angelo, *Thin Solid Films* **603** (2016) 149-153.
12. 時間分解軟X線光電子分光法：半導体表面における光励起キャリアの実時間観測：山本 達，松田 巖，表面科学 **37** (2016) 9-13.
13. Capturing transient charged states at the C₆₀/TiO₂(110) interface by time-resolved soft X-ray photoelectron spectroscopy: K. Ozawa, S. Yamamoto, R. Yukawa, K. Akikubo, M. Emori, H. Sakama and I. Matsuda, *Organic Electronics* **31** (2016) 98-103.
14. * Real-time observation of reaction processes of CO₂ on Cu(997) by ambient-pressure X-ray photoelectron spectroscopy: T. Koitaya, S. Yamamoto, Y. Shiozawa, K. Takeuchi, R.-Y. Liu, K. Mukai, S. Yoshimoto, K. Akikubo, I. Matsuda and J. Yoshinobu, *Topic in Catalysis* **59** (2016) 526-531.

Kobayashi group

We have demonstrated the highest repetition rate in a Kerr-lens mode-locked oscillator in the world. 15-GHz mode space of the optical frequency comb can be resolved by using commercially available spectrum analyzer. We started to develop a new HHG beam line for industrial applications. We have demonstrated 300-mW, 193-nm light source with single longitudinal mode at 6 kHz repetition rate for next generation lithography technology.

1. †300-mW narrow-linewidth deep-ultraviolet light generation at 193 nm by frequency mixing between Yb-hybrid and Er-fiber lasers: H. Xuan, Z. Zhao, H. Igarashi, S. Ito, K. Kakizaki and Y. Kobayashi, *Opt. Express* **23** (2015) 10564-10572.
2. Direct 15-GHz mode-spacing optical frequency comb with a Kerr-lens mode-locked Yb:Y₂O₃ ceramic laser: M. Endo, I. Ito and Y. Kobayashi, *Opt. Express* **23** (2015) 1276-1282.
3. †High average power coherent vuv generation at 10 MHz repetition frequency by intracavity high harmonic generation: A. Ozawa, Z. Zhao, M. Kuwata-Gonokami and Y. Kobayashi, *Opt. Express* **23** (2015) 15107-15118.
4. Offset-free broadband Yb: fiber optical frequency comb for optical clocks: T. Nakamura, I. Ito and Y. Kobayashi, *Opt. Express* **23** (2015) 19376-19381.
5. †Single frequency, 5 ns, 200 μJ, 1553 nm fiber laser using silica based Er-doped fiber: Z. Zhao, H. Xuan, H. Igarashi, S. Ito, K. Kakizaki and Y. Kobayashi, *Opt. Express* **23** (2015) 29764-29771.
6. †* Emergent photovoltage on SmB₆ surface upon bulk-gap evolution revealed by pump-and-probe photoemission spectroscopy: Y. Ishida, T. Otsu, T. Shimada, M. Okawa, Y. Kobayashi, F. Iga, Y. Takabatake and S. Shin, *Sci. Rep.* **5** (2015) 8160(1-6).
7. †Femtosecond optical parametric oscillator frequency combs: Y. Kobayashi, K. Torizuka, A. Marandi, R. L. Byer, R. A. McCracken, Z. Zhang and D. T. Reid, *J. Opt.* **17** (2015) 094010(1-14).
8. Ytterbium fiber-based, 270 fs, 100 W chirped pulse amplification laser system with 1 MHz repetition rate: Z. Zhao and Y. Kobayashi, *Appl. Phys. Express* **9** (2016) 012701(1-4).
9. * High-precision group-delay dispersion measurements of optical fibers via fingerprint-spectral wavelength-to-time mapping: T. Ito, O. Slezak, M. Yoshita, H. Akiyama and Y. Kobayashi, *Photon. Res.* **4** (2016) 13-16.
10. 5. レーザー (2014 年日本光学会の研究動向)：赤羽 浩一，小林 洋平，光学 **44** (2015) 141-144.
11. ファイバレーザ光周波数コム技術と光科学への展開：小林 洋平，月刊オプトロニクス **34** (2015) 144-147.

* Joint research among groups within ISSP.

Itatani group

We carried out several application experiments using the optical parametric chirped pulse amplification (OPCPA) system that can produce carrier-envelope phase (CEP)-stable, 1.5-mJ, 10-fs infrared pulses at a repetition rate of 1 kHz. First, we produced soft-X-ray continuum extending to the photon energy of 320 eV, and measured the static absorption spectrum of a thin film containing carbon atoms. We successfully resolved the peaks arising from the C=C and C-C bonding with an accumulation time of approximately 100 seconds. This result shows the capability of laser-based time-resolved soft-X-ray absorption spectroscopy. Second, we measured the photoelectron spectra of ionizing atoms, and observed the CEP-dependent high-energy structures. By mapping the CEP-dependent cutoffs of photoelectron spectra, we successfully reproduced the differential cross sections of rare gas atoms with a high degree of agreement with the most advanced scattering theory. Third, under the collaboration with Prof. Tanaka at Kyoto University, we produced broadband THz pulses using organic crystals. New directions of the BIBO-based ultrabroadband optical parametric amplifiers are pursuit as well. First, we developed a high-repetition-rate optical parametric amplifier using newly developed dispersion compensation mirrors. We have successfully produced CEP-stable 10 uJ, 9-fs infrared pulses at 20 kHz, which were applied to high-throughput electron scattering experiments. Second, we modified a high-energy OPCPA system to amplify two spectral components in infrared followed by differential frequency generation, resulting in the generation of intense mid-infrared pulses. A strong optical field up to 50 MV/cm was achieved with a capability of sub-cycle EO sampling using 6.5-fs visible pulses. We also kept collaboration with Shin, Komori, and Matsuda groups at LASOR-ISSP on time-resolved ARPES using a femtosecond EUV source. Photo-induced electronic dynamics of $\text{Cu}_{0.17}\text{Bi}_2\text{Se}_3$ and graphene are successfully measured.

1. ^{†*}Nonequilibrium electronic and phonon dynamics of $\text{Cu}_{0.17}\text{Bi}_2\text{Se}_3$ investigated by core-level and valence-band time-resolved photoemission spectroscopy: T. Yamamoto, Y. Ishida, R. Yoshida, M. Okawa, K. Okazaki, T. Kanai, A. Kikkawa, Y. Taguchi, T. Kiss, K. Ishizaka, N. Ishii, J. Itatani, S. Watanabe, Y. Tokura, T. Saitoh and S. Shin, *Phys. Rev. B* **92** (2015) 121106(1-6).
2. Ultrabroadband infrared chirped mirrors characterized by a white-light Michelson interferometer: K. Kaneshima, M. Sugiura, K. Tamura, N. Ishii and J. Itatani, *Appl. Phys. B* **119** (2015) 347-353.
3. ^{†*}フェムト秒域時間分解光電子分光法によるグラフェンの超高速キャリアダイナミクスの追跡 Tracing Ultrafast Carrier Dynamics in Graphene with Femtosecond Time-resolved Photoemission Spectroscopy: 染谷 隆史, 吹留 博一, 石田 行章, 吉田 力矢, 山本 達, 板谷 治郎, 小森 文夫, 辛 埴, 松田 巖, *表面科学* **36(8)** (2015) 418-423.
4. [†]Generation of ultrashort intense optical pulses at 1.6 μm from a bismuth triborate-based optical parametric chirped pulse amplifier with carrier-envelope phase stabilization: N. Ishii, K. Kaneshima, T. Kanai, S. Watanabe and J. Itatani, *J. Optics* **17** (2015) 094001(1-8).
5. Generation of carrier-envelope phase-stable mid-infrared pulses via dual-wavelength optical parametric amplification: K. Kaneshima, N. Ishii, K. Takeuchi and J. Itatani, *Opt. Express* **24** (2016) 8660(1-6).

Harada group

After three years development of differential pumping system we have for the first time succeeded in the near ambient pressure soft X-ray RIXS experiments at SPring-8 BL07LSU. The best demonstration was done for O 1s RIXS of SiO_2 under around 360 Torr (0.5 bar) air pressure. We also have implemented rotation of the RIXS spectrometer to enable momentum dependent RIXS experiments urgently requested for the study of strongly correlated systems. This year we have accepted 8 collaborative works at BL07LSU HORNET endstation, which include oxygen site analysis of high T_c cuprates, operando analysis of Li ion battery electrodes, RIXS study of multiferroic materials applying magnetic field, spin transition of LaCoO_3 , monitoring vibrational excitations at Ti site in ferroelectric BaTiO_3 , and electronic structure analysis of high concentration electrolyte for Li ion batteries.

1. [†]Multi-Phonon Excitations in Fe 2p RIXS on Mg_2FeH_6 : K. Kurita, D. Sekiba, I. Harayama, K. Chito, Y. Harada, H. Kiuchi, M. Oshima, S. Takagi, M. Matsuo, R. Sato, K. Aoki and S.-I. Orimo, *J. Phys. Soc. Jpn.* **84** (2015) 043201(1-3).
2. [†]Electron-Phonon Coupling in the Bulk of Anatase TiO_2 Measured by Resonant Inelastic X-Ray Spectroscopy: S. Moser, S. Fatale, P. Krüger, H. Berger, P. Bugnon, A. Magrez, H. Niwa, J. Miyawaki, Y. Harada and M. Grioni, *Phys. Rev. Lett.* **115** (2015) 096404 (1-5).
3. X-ray Emission Spectroscopy of Bulk Liquid Water in "No-man's Land": J. Sellberg, T. McQueen, H. Laksmono, S. Schreck, M. Beye, D. DePonte, B. O'Kennedy, D. Nordlund, R. Sierra, D. Schlesinger, T. Tokushima, S. Eckert, V. Segtnan, H. Ogasawara, K. Kubicek, S. Techert, U. Bergmann, G. Dakovski, W. Schlotter, Y. Harada, I. Zhovtobriukh, M. Bogan, P. Wernet, A. Föhlisch, L. Pettersson and A. Nilsson, *J. Chem. Phys.* **142** (2015) 044505(1-9).
4. [†]Enhancement in Kinetics of the Oxygen Reduction Reaction on a Nitrogen-Doped Carbon Catalyst by Introduction of Iron via Electrochemical Methods: J. Wu, D. Zhang, H. Niwa, Y. Harada, M. Oshima, H. Ofuchi, Y. Nabae, T. Okajima and T. Ohsaka, *Langmuir* **31** (2015) 5529-5536.

[†] Joint research with outside partners.

5. Active site formation mechanism of carbon-based oxygen reduction catalysts derived from a hyperbranched iron phthalocyanine polymer: Y. Hiraike, M. Saito, H. Niwa, M. Kobayashi, Y. Harada, M. Oshima, J. Kim, Y. Nabae and M.-A. Kakimoto, *Nanoscale Res Lett* **10** (2015) 179(1-11).
6. [†]Operando soft x-ray emission spectroscopy of LiMn₂O₄ thin film involving Li-ion extraction/insertion reaction: D. Asakura, E. Hosono, H. Niwa, H. Kiuchi, J. Miyawaki, Y. Nanba, M. Okubo, H. Matsuda, H. S. Zhou, M. Oshima and Y. Harada, *Electrochem. Commun.* **50** (2015) 93-96.
7. ^{*}First-Principles Investigation of Strong Excitonic Effects in Oxygen 1s X-ray Absorption Spectra: Y. Noguchi, M. Hiyama, H. Akiyama, Y. Harada and N. Koga, *J. Chem. Theory Comput.* **11** (2015) 1668-1673.
8. [†]Redox Potential Paradox in Na_xMO₂ for Sodium-Ion Battery Cathodes: Y. Nanba, T. Iwao, B. M. D. Boisse, W. Zhao, E. Hosono, D. Asakura, H. Niwa, H. Kiuchi, J. Miyawaki, Y. Harada, M. Okubo and A. Yamada, *Chem. Mater.* **28** (2016) 1058-1065.
9. In Situ Hard X-ray Photoelectron Study of O₂ and H₂O Adsorption on Pt Nanoparticles: Y. Cui, Y. Harada, E. Ikenaga, R. Li, N. Nakamura, T. Hatanaka, M. Ando, T. Yoshida, G.-L. Li and M. Oshima, *J. Phys. Chem. C* **120** (2016) 10936-10940.
10. [†]Characterization of nitrogen species incorporated into graphite using low energy nitrogen ion sputtering: H. Kiuchi, T. Kondo, M. Sakurai, D. Guo, J. Nakamura, H. Niwa, J. Miyawaki, M. Kawai, M. Oshima and Y. Harada, *Phys. Chem. Chem. Phys.* **18** (2016) 458-465.
11. Pt-free carbon-based fuel cell catalyst prepared from spherical polyimide for enhanced oxygen diffusion: Y. Nabae, S. Nagata, T. Hayakawa, H. Niwa, Y. Harada, M. Oshima, A. Isoda, A. Matsunaga, K. Tanaka and T. Aoki, *Sci. Rep.* **6** (2016) 23276(1-7).
12. [†]Lewis Basicity of Nitrogen-Doped Graphite Observed by CO₂ Chemisorption: H. Kiuchi, R. Shibuya, T. Kondo, J. Nakamura, H. Niwa, J. Miyawaki, M. Kawai, M. Oshima and Y. Harada, *Nanoscale Res Lett* **11** (2016) 127(1-7).
13. [†]Intermediate honeycomb ordering to trigger oxygen redox chemistry in layered battery electrode: B. M. D. Boisse, G. Liu, J. Ma, S.-I. Nishimura, S.-C. Chung, H. Kiuchi, Y. Harada, J. Kikkawa, Y. Kobayashi, M. Okubo and A. Yamada, *Nat. Commun.* **7** (2016) 11397(1-9).

Wadati group

We succeeded in the observation of a devil's staircase in the novel spin-valve system SrCo₆O₁₁ by resonant soft x-ray scattering. We also performed time-resolved reflectivity study of ferrimagnetic alloy GdFeCo thin films by using seeded free electron laser in FERMI (Italy), and observed ultrafast spin-switching of Fe spins by resonant magneto-optical Kerr effect.

1. Investigating Orbital Magnetic Moments in Spinel-Type MnV₂O₄ Using X-ray Magnetic Circular Dichroism: J. Okabayashi, S. Miyasaka, K. Hemmi, K. Tanaka, S. Tajima, H. Wadati, A. Tanaka, Y. Takagi and T. Yokoyama, *J. Phys. Soc. Jpn.* **84** (2015) 104703(1-5).
2. X-ray spectroscopic study of BaFeO₃ thin films; an Fe⁴⁺ ferromagnetic insulator: T. Tsuyama, T. Matsuda, S. Chakraborty, J. Okamoto, E. Ikenaga, A. Tanaka, T. Mizokawa, H. Y. Hwang, Y. Tokura and H. Wadati, *Phys. Rev. B* **91** (2015) 115101(1-7).
3. ^{*}Ultrafast spin-switching of a ferrimagnetic alloy at room temperature traced by resonant magneto-optical Kerr effect using a seeded free electron laser: Sh. Yamamoto, M. Taguchi, T. Someya, Y. Kubota, S. Ito, H. Wadati, M. Fujisawa, F. Capotondi, E. Pedersoli, M. Manfredda, L. Raimondi, M. Kiskinova, J. Fujii, P. Moras, T. Tsuyama, T. Nakamura, T. Kato, T. Higashide, S. Iwata, S. Yamamoto, S. Shin and I. Matsuda, *Rev. Sci. Instrum.* **86** (2015) 083901(1-5).
4. Engineering a spin-orbital magnetic insulator by tailoring superlattices: J. Matsuno, K. Ihara, S. Yamamura, H. Wadati, K. Ishii, V. V. Shankar, H.-Y. Kee and H. Takagi, *Phys. Rev. Lett.* **114** (2015) 247209(1-5).
5. Observation of a Devil's Staircase in the Novel Spin-Valve System SrCo₆O₁₁: T. Matsuda, S. Partzsch, T. Tsuyama, E. Schierle, E. Weschke, J. Geck, T. Saito, S. Ishiwata, Y. Tokura and H. Wadati, *Phys. Rev. Lett.* **114** (2015) 236403(1-5).
6. Electronic structure of Li_{1+x}[Mn_{0.5}Ni_{0.5}]_{1-x}O₂ studied by photoemission and x-ray absorption spectroscopy: Y. Yokoyama, D. Ootsuki, T. Sugimoto, H. Wadati, J. Okabayashi, X. Yang, F. Du, G. Chen and T. Mizokawa, *Appl. Phys. Lett.* **107** (2015) 033903(1-4).
7. Topotactic synthesis of strontium cobalt oxyhydride thin film with perovskite structure: T. Katayama, A. Chikamatsu, H. Kamisaka, Y. Yokoyama, Y. Hirata, H. Wadati, T. Fukumura and T. Hasegawa, *AIP Advances* **5** (2015) 107147(1-7).

^{*} Joint research among groups within ISSP.

8. 新型スピントロニクスと悪魔の階段：和達 大樹，パリティ **31** (2016) 48-51.
9. Material/element-dependent fluorescence-yield modes on soft X-ray absorption spectroscopy of cathode materials for Li-ion batteries: D. Asakura, E. Hosono, Y. Nanba, H. Zhou, J. Okabayashi, C. Ban, P. -A. Glans, J. Guo, T. Mizokawa, G. Chen, A. J. Achkar, D. G. Hawthorn, T. Z. Regier and H. Wadati, AIP Advances **6** (2016) 035105(1-8).

Kondo group

We use angle-resolved photoemission spectroscopy (ARPES) with ultrahigh energy resolution. The main findings in 2015 are as follows: (1) Quadratic Fermi Node in a 3D Strongly Correlated Semimetal. (2) Point nodes persisting far beyond T_c in Bi2212.

1. Tunable spin current due to bulk insulating property in the topological insulator $Tl_{1-x}Bi_{1+x}Se_{2-\delta}$: K. Kuroda, G. Eguchi, K. Shirai, M. Shiraishi, M. Ye, K. Miyamoto, T. Okuda, S. Ueda, M. Arita, H. Namatame, M. Taniguchi, Y. Ueda and A. Kimura, Phys. Rev. B **91** (2015) 205306.
2. ^{†*}Point nodes persisting far beyond T_c in Bi2212: T. Kondo, W. Malaeb, Y. Ishida, T. Sasagawa, H. Sakamoto, T. Takeuchi, T. Tohyama and S. Shin, Nat. Commun. **6** (2015) 7699(8 pages).
3. ^{†*}Quadratic Fermi Node in a 3D Strongly Correlated Semimetal: T. Kondo, M. Nakayama, R. Chen, J. J. Ishikawa, E. -G. Moon, T. Yamamoto, Y. Ota, W. Malaeb, H. Kanai, Y. Nakashima, Y. Ishida, R. Yoshida, H. Yamamoto, M. Matsunami, S. Kimura, N. Inami, K. Ono, H. Kumigashira, S. Nakatsuji, L. Balents and S. Shin, Nat. Commun. **6** (2015) 10042(1-8).
4. ^{*}Optically detecting the edge-state of a three-dimensional topological insulator under ambient conditions by ultrafast infrared photoluminescence spectroscopy: S.-Y. Maezawa, H. Watanabe, M. Takeda, K. Kuroda, T. Someya, I. Matsuda and T. Suemoto, Sci. Rep. **5** (2015) 16443(1-8).
5. ^{†*}Ultrafast electron dynamics at the Dirac node of the topological insulator Sb_2Te_3 : S. Zhu, Y. Ishida, K. Kuroda, K. Sumida, M. Ye, J. Wang, H. Pan, M. Taniguchi, S. Qiao, S. Shin and A. Kimura, Sci. Rep. **5** (2015) 13213(1-6).
6. Pairing, pseudogap and Fermi arcs in cuprates.: A. Kaminski, T. Kondo, T. Takeuchi and G. Gu, Philosophical Magazine **95** (2015) 453-466.
7. ^{*}High-resolution three-dimensional spin- and angle-resolved photoelectron spectrometer using vacuum ultraviolet laser light: K. Yaji, A. Harasawa, K. Kuroda, S. Toyohisa, M. Nakayama, Y. Ishida, A. Fukushima, S. Watanabe, C. Chen, F. Komori and S. Shin, Rev. Sci. Instrum. **87** (2016) 053111(1-6).
8. Generation of Transient Photocurrents in the Topological Surface State of Sb_2Te_3 by Direct Optical Excitation with Midinfrared Pulses: K. Kuroda, J. Reimann, J. Gdde and U. Hfer, Phys. Rev. Lett. **116** (2016) 076801.
9. ^{†*}Spin Polarization and Texture of the Fermi Arcs in the Weyl Fermion Semimetal TaAs: S.-Y. Xu, I. Belopolski, D. S. Sanchez, M. Neupane, G. Chang, K. Yaji, Z. Yuan, C. Zhang, K. Kuroda, G. Bian, C. Guo, H. Lu, T.-R. Chang, N. Alidoust, H. Zheng, C.-C. Lee, S.-M. Huang, C.-H. Hsu, H.-T. Jeng, A. Bansil, T. Neupert, F. Komori, T. Kondo, S. Shin, H. Lin, S. Jia and M. Zahid Hasan, Phys. Rev. Lett. **116** (2016) 096801(1-7).
10. ^{†*}角度分解光電子分光による精密測定で解き明かす銅酸化物高温超伝導体の擬ギャップと超伝導ギャップの競合関係：近藤 猛，竹内 恒博，辛 埴，固体物理 **51** (2016) 203-221.

Okazaki group

We have developed and improved a time- and angle-resolved photoemission (TrARPES) apparatus using EUV and SX lasers by high harmonics generation (HHG). In the fiscal year 2015, we have installed a new Ti:Sapphire regenerative amplifier system and fairly improved a performance and stability of this apparatus. We have studied nonequilibrium electronic states of semiconductor surfaces, graphene, high- T_c cuprates, iron-based superconductors, and strongly correlated electron systems. Particularly, we have observed coherent phonon excitations both in the hole and electron Fermi surfaces of a parent compound of iron-based superconductors $BaFe_2As_2$. In addition, we have also investigated superconducting-gap structures of iron-based superconductors and BiS2-based superconductors by a low-temperature and high-resolution laser ARPES apparatus.

1. ^{†*}Low-Temperature and High-Energy-Resolution Laser Photoemission Spectroscopy: T. Shimojima, K. Okazaki and S. Shin, J. Phys. Soc. Jpn. **84** (2015) 072001(1-26).
2. Dependence of electron correlation strength in $Fe_{1+y}Te_{1-x}Se_x$ on Se content: L. C. C. Ambolode, K. Okazaki, M. Horio, H. Suzuki, L. Liu, S. Ideta, T. Yoshida, T. Mikami, T. Kakeshita, S. Uchida, K. Ono, H. Kumigashira, M. Hashimoto, D. -H. Lu, Z. -X. Shen and A. Fujimori, Phys. Rev. B **92** (2015) 035104.

[†] Joint research with outside partners.

3. In-plane electronic anisotropy in the antiferromagnetic orthorhombic phase of isovalent-substituted $\text{Ba}(\text{Fe}_{1-x}\text{Ru}_x)_2\text{As}_2$: L. Liu, T. Mikami, S. Ishida, K. Koshiishi, K. Okazaki, T. Yoshida, H. Suzuki, M. Horio, L. C. C. Ambolode, J. Xu, H. Kumigashira, K. Ono, M. Nakajima, K. Kihou, C. H. Lee, A. Iyo, H. Eisaki, T. Kakeshita, S. Uchida and A. Fujimori, *Phys. Rev. B* **92** (2015) 094503(1-5).
4. ^{†*}Nonequilibrium electronic and phonon dynamics of $\text{Cu}_{0.17}\text{Bi}_2\text{Se}_3$ investigated by core-level and valence-band time-resolved photoemission spectroscopy: T. Yamamoto, Y. Ishida, R. Yoshida, M. Okawa, K. Okazaki, T. Kanai, A. Kikkawa, Y. Taguchi, T. Kiss, K. Ishizaka, N. Ishii, J. Itatani, S. Watanabe, Y. Tokura, T. Saitoh and S. Shin, *Phys. Rev. B* **92** (2015) 121106(1-6).
5. Reversed anisotropy of the in-plane resistivity in the antiferromagnetic phase of iron tellurides: L. Liu, T. Mikami, M. Takahashi, S. Ishida, T. Kakeshita, K. Okazaki, A. Fujimori and S. Uchida, *Phys. Rev. B* **91** (2015) 134502.
6. ^{†*} 高次高調波の固体時間分解光電子分光への応用 Application of High Harmonic Generation to Time-Resolved Photoemission Spectroscopy of Solids: 岡崎 浩三, 辛 埴, レーザー研究 **43** (12) (2015) 838-843.
7. Suppression of the antiferromagnetic pseudogap in the electron-doped high-temperature superconductor by protect annealing: M. Horio, T. Adachi, Y. Mori, A. Takahashi, T. Yoshida, H. Suzuki, L. C. C. Ambolode, K. Okazaki, K. Ono, H. Kumigashira, H. Anzai, M. Arita, H. Namatame, M. Taniguchi, D. Ootsuki, K. Sawada, M. Takahashi, T. Mizokawa, Y. Koike and A. Fujimori, *Nat. Commun.* **7** (2016) 10567(1-8).

* Joint research among groups within ISSP.